

SOV/49-58-7-6/16

Statistical Methods of Determination of Effective Parameters of
the Observed Convective Currents

The concentration of N_1 , i.e. the number of circles per unit of surface, can be expressed by the Eq.(10) where n is the number of currents crossed by an aircraft taking temperatures along the path L .

b) In the case of air bubbles, apart from their shape, it is necessary to consider the question of their spatial distribution. This can be expressed as a distribution of the ellipsoids (11) and (12) with their concentration. The probability of horizontal cross-sections of the ellipsoids having the dimensions from l to $l + dl$ will be Eq.(13). The conditional probability (14) is equal to the ratio of the elliptic rings (having axes a_1 and a_1/m with thickness da_1) to the surface of the ellipse (having axes a and a/m). If Eq.(15) is the density probability function of the ellipsoids distribution and N_2 the number of the ellipsoid centres per unit of the volume, the equation (16) is the number of ellipsoids along the horizontal

Card3/11 straight line (a^2 = the second moment). From Eqs.(17) and

SOV/49-58-7-6/16

Statistical Methods of Determination of Effective Parameters of
the Observed Convective Currents

(14) substituted into Eq.(13), an integral equation (18) can be formed which expresses the relationship between the distribution of the ellipsoids and the chords. The solution for this equation is given by (19).

The concentration of ellipsoids N_0 , i.e. their number per unit of volume, according to Eq.(16) can be expressed as Eq.(20), where n = total number of ellipsoids along the path L .

It should be noted that the Eq.(18) contains no reference to m . Therefore, it can be applied to all shapes of the air bubbles, such as sphere ($m = 1$), vertical ellipsoid ($m < 1$) or horizontal ellipsoid ($m > 1$).

3) A relative surface S of the air current at any cross-section is expressed by Eq.(21) where \bar{s}^2 = second moment.

The value of S can be obtained from the observations.

By multiplying both parts of Eq.(8) by l and integrating for l , the Eq.(22) is obtained (\bar{l} = mean chord).

The number of ellipsoids (23) on a horizontal surface (cross-section) can be calculated from Eqs.(3) and (5) where

Card4/11

SOV/49-58-7-6/16

Statistical Methods of Determination of Effective Parameters of the Observed Convective Currents

\bar{a} = mean length of the horizontal axes. The density of ellipsoids intersecting a surface $u(a)$ is Eq.(24). The probability of obtaining the circles of diameters from s to $s + ds$ on the intersected surface is Eq.(25) and the conditional probability of the surface being formed by cutting an ellipsoid at the distance from its centre y to $y + dy$ will be Eq.(26). From Eqs.(26) and (24) substituted into (25), the equation (27), expressing the distribution of circles on a horizontal surface is formed. It is assumed that a surface under consideration represents the surface of flight. Then, taking Eq.(28) instead of $F_2(a)$ in Eq.(19) and $N_1 = N_2 \bar{a}/m$ from Eq.(23), an equation (29) is obtained. This is the same as (9), thus (30) is being formed, which indicates that the relative surface of the ascending air in both cases can be determined by Eqs.(21) or (22). Also, it is evident from Eq.(30) that the distribution of circles on a horizontal surface is independent from the kind of air convection, The relative volume V of the air bubbles can also be

Card5/11

SOV/49-58-7-6/16

Statistical Methods of Determination of Effective Parameters of the Observed Convective Currents

shown as Eq.(31), where \bar{a}^3 = third moment. In this case, the relative volume of the convective current is equal to the relative surface of any horizontal cross-section.

4) The equations can be applied in the cases of air being either stationary or in motion with winds gradually varying with height. In the latter case, the convective currents will be of an inclined elliptic shape. But their cross-section will be represented as a circle. Therefore, the Eqs.(9) (10) and (21) (22) can be applied. Similarly, Eqs.(14) and (19)(20) can also be applied for the air bubble in these conditions.

5) In order to determine the statistical relationship between the dimension and the temperature of the convective current, it is necessary to consider their joint distribution, i.e. to solve a question with the two independent variables.

In the case of air bubbles, the probability of the horizontal cross-section of ellipsoid having a given rate of the temperature increase is calculated by Eq.(32).

Card6/11

SOV/49-58-7-6/16

Statistical Methods of Determination of Effective Parameters of the Observed Convective Currents

This equation can only be solved when the temperature condition (33) is satisfied. Therefore, Eq.(34) will be integrated in respect of one variable only. If the internal temperature of the ellipsoid changes according to Eq.(36), then Eq.(37) will be obtained. Here, λ and T are two independent variables with T included in the integral as a parameter. By introducing a new variable, c (38), the Eq.(37) will change into (39), where c can be considered as a parameter. After both parts being differentiated in respect of λ , the function (40) or (41) is obtained.

In the case of the airstream, the temperature change will follow Eq.(42) and the above probability will be expressed by (43), which can be written as (44), where $T_{0,s}$ is

the internal temperature varying according to Eq.(45). By the inclusion of Eq.(45) into F_2 in Eq.(44), the form (46) is obtained, which can be written as Eq.(47).

Assuming that the profile of the temperature distribution (48) represents a circle, the Eqs.(49) and (50) will be

Card7/11

Statistical Methods of Determination of Effective Parameters of the
Observed Convective Currents

SOV/49-58-7-6/16

obtained. These are solved by substituting the variable by $T = c_l$ in Eq.(51) in order to obtain the final solutions (52) and (53).

6) If more variables are included which could represent a number of conditions in the interior of ellipsoids or circles, the Eqs.(54) to (59) will be necessary. The previously derived Eqs.(8), (18), (37), (43), (49), (50 and their solutions (9), (19), (40), (46), (52), (53) can be treated as the particular cases of (54) and (55) with the solutions (58) and (59).

7) As it was shown before, it is possible to consider the distribution of circles or ellipsoids grouped according to their dimensions and the central temperatures instead of the distribution of chords. In order to do that, Eqs.(60) and (61), formed from (8) and (18), should be introduced, then Eq.(62) will be found. In the case of two independent variables, as it may occur when the temperature changes according to Eqs.(36) or (42), the required distribution will follow the Eqs.(63) to (66). Taking the

Card8/11

SOV/49-58-7-6/16

Statistical Methods of Determination of Effective Parameters of
the Observed Convective Currents

temperature changes (48) instead of (63) and (64), the Eqs.(67) and (68) will be derived with the limits similar to Eqs.(65) and (66).
8) The Eqs.(9),(19),(40),(46),(52) and (53) express the distribution based on the functions $w(l)$ or $w(l, T)$ found experimentally from a great number of flights. However, a simpler method can be applied for calculation of two or more independent variables.
Eqs.(37) and (43) with the substitution (38) or Eqs.(49) and (50) with (51) are transformed independently of the variables l and c . In particular, the Eq.(43) will be obtained in a form (69). Denoting the number of the simultaneous observations as $f_{k+0.5; i+0.5}$, the Equations (70) and (71) are formed. By changing the succession of integration of the last two integrals and by integrating in respect of l , the Eq.(72) will be obtained. The second integral can be presented in the form of Eq.(73) and instead of (72), a system of algebraic equations (74) to (76) can be used with the coefficients being independent

Card9/11

SOV/49-58-7-6/16

Statistical Methods of Determination of Effective Parameters of
the Observed Convective Currents

of k . If Eqs.(49) and (50) are included, these equations become (77) and (78).

The equations of the type (74) are easy to evaluate. The evaluation of (74) and (75) with the coefficients is shown in Tables 1 and 2. The tables show the distribution of dimensions at an interval c which was taken from the Eqs.(79) and (80). The data from the tables can be considered as having only one independent variable, i.e. the index $k+0.5$ being omitted.

9) As an example, the data of 5 flights over the Ukraine in June, 1956, is given. The total time of the flights was 10 - 13 hours at heights ranging from 10 - 50 m to 1 500 - 2 000 m. The majority of observations were made at 300 - 500 m (Table 2).

Figure 1 shows the probability density functions of the following distributions: chord dimension $w(l)$, circle dia. $F_1(s)$, horizontal axis of ellipsoid $F_2(a)$.

Figure 2 represents the probability density functions of two independent variables of the following distributions:

Card10/11

SOV/49-58-7-6/16
Statistical Methods of Determination of Effective Parameters of
the Observed Convective Currents

chord dimension and temperature $w(l, T)$ (the upper graph),
circle diameter and temperature $F_1(s, T_0)$ (the middle
graph), horizontal axis of ellipsoid and temperature
 $F_2(a, T_0)$ (the lower graph).

There are 2 figures and 2 tables and 9 references,
5 of which are Soviet, 3 English and 1 French.

ASSOCIATION: Akademiya nauk SSSR Institut prikladnoy geofiziki
(Institute of Applied Geophysics, Ac.Sc.USSR)

SUBMITTED: December 23, 1957

Card 11/11

- | | |
|-------------------------|------------------------------|
| 1. Atmosphere--Motion | 2. Clouds--Development |
| 3. Convection--Analysis | 4. Mathematics--Applications |

VUL'FSON, N.I., Doc Phys-Math Sci -- (diss) "Study of convective
^{movements} motions in ^{the} free atmosphere." Mos, 1959. 14 pp (Acad Sci USSR. Inst
of Applied Geophysics). 225 copies. List of author's works at
end of text (10 titles) (KL, 37-59, 105)

3 (8)

AUTHOR:

Vul'fson, N. I.

SOV/20-126-6-27/67

TITLE:

On the Mechanism of ^{Permissible} Instability ~~Loss~~ in the Free Atmosphere (O mekhanizme razresheniya neustoychivosti v svobodnoy atmosfere)

PERIODICAL:

Doklady Akademii nauk SSSR, 1959, Vol 126, Nr 6, pp 1244 - 1247 (USSR)

ABSTRACT:

In recent years a method was developed of announcing convective rising airflows by the aid of sensitive thermometers. The temperature distribution in the flow is determined by statistical integration of the measuring results. The distribution function (1) and the temperature profile (3) of the flow are given. Measurements reveal that the temperature distribution profile varies but little with altitude; figure 1 shows a graph depicting the variation of temperature in the center of the flow with altitude. Formula (4) describes the temperature drop in the axis of flow. Next, equations (5) and (6) are given for the temperature- and velocity variations in the axis of turbulent flows. These equations are expanded for any arbitrary point in the flow (9) (10). Results reveal that dimensions of convective flow systematically decrease with rising instability and turbu-

Card 1/2

On the Mechanism of Evacuation of Instability Lease SOV/20-126-6-27/67
the Free Atmosphere

lence of atmosphere. Furthermore, it follows that instability evacuates in free atmosphere in the form of a spontaneous convective flow. There are 2 figures, 2 tables, and 9 references, 5 of which are Soviet.

ASSOCIATION: Institut prikladnoy geofiziki Akademii nauk SSSR (Institute of Applied Geophysics of the Academy of Sciences, USSR)

PRESENTED: February 16, 1959, by L. I. Sedov, Academician

SUBMITTED: February 11, 1959

Card 2/2

PHASE I BOOK EXPLOITATION SOV/5682

Vul'fson, Naum Isaakovich

Issledovaniye konvektivnykh dvizheniy v svobodnoy atmosfere .
(Investigation of the Convective Motions in the Free Atmos-
phere) Moscow, Izd-vo AN SSSR, 1961. 251 p. Errata slip
inserted. 1,500 copies printed.

Sponsoring Agency: Akademiya nauk SSSR. Institut prikladnoy
geofiziki.

Resp. Ed.: I. A. Kibel', Corresponding Member, Academy of Sciences
USSR; Ed. of Publishing House: G. G. Gus'kov; Tech. Eds.: T. A.
Prusakova and G. N. Shevchenko.

PURPOSE : This book is intended for scientific workers in physics
of the atmosphere and certain related fields (e.g., aviation and
atmospheric pollution control). It may also be used by students
in meteorological institutes and faculty members of university
departments of physics.

Card ~~1/7~~ 1/3

Investigation of the Convective (Cont.)

SOV/5682

COVERAGE: The book presents results of an investigation of convective motions in the free atmosphere and in cumulus clouds. A large portion of the book is devoted to substantiating the observational procedures and instrumentation of aircraft flights required for acquiring the basic data, and methods of processing, studying, and statistically interpreting the observational results. Basic convection parameters and their variation with changing physiogeographical conditions, the mechanism for the development of convective motions in the free atmosphere, and the association between convective motions within and without clouds can be established from the data presented. The obtained results are valuable for understanding the mechanism of convection and the development of convective clouds and for their possible uses as quantitative indices of phenomena associated with convective motions. The author thanks Academician Ye. K. Fedorov (scientific advice); V. V. Shchelokov, V. I. Skatskiy, A. M. Gromov (design of apparatus); N. V. Davydkin and I. S. Pavlova (installation and operation of apparatus); A. I. Korzhov, V. N. Shlyakov (deceased), P. N. Radkevich

Card 2/13

Investigation of the Convective (Cont.)

SOV/5682

and N. S. Kolesov (commanders of flight crews); V. M. Bovsheverov, L. M. Levin, I. A. Kibel', Corresponding Member AS USSR, A. M. Obukhov, Corresponding Member AS USSR, A. Kh. Khrgian, Professor, and N. Z. Pinus, Doctor of Physics and Mathematics (critical remarks); S. V. Pshenay-Severin (editorial assistance). There are 149 references: 61 Soviet, 70 English, 9 German, 8 French, and 1 Polish.

TABLE OF CONTENTS:

Author's Foreword	3
Introduction	5
Ch. I. Method of Studying Convective Motions in the Free Atmosphere	15
1. Physical bases of the method	15
2. Thermometric assembly on aircraft	21
3. Errors of measurements	33

Card 3/7₃

VUL'FSON, N.I., doktor fiz.-matem. nauk, otv. red.; LEVIN, L.M.,
doktor fiz.-matem. nauk, otv. red. Prinimali uchastiye:
KOMAROV, N.H., red.; PSHENAY-SEVERIN, S.V., red.; UGAROVA, K.F.,
red.; NIKOLAYEVA, L.K., red. izd-va; BERKGAUT, V.G., red. izd-va;
VOLKOVA, V.V., tekhn. red.

[Study of clouds, precipitation, and thunderstorm electricity;
reports] Issledovaniia oblakov, osadkov i grozovogo elektri-
chestva; doklady. Otv. red. N.I.Vul'fson, L.M.Levin. Moskva,
Izd-vo Akad.nauk SSSR. 1961. 327 p. (MIRA 15:1)

1. Mezhdomstvennaya konferentsiya po voprosam issledovaniya
oblakov, osadkov i atmosfernogo elektrichestva. 6th, 1959.
(Cloud physics—Congresses)

VUL'FSON, N.I.

Irregularities of the atmospheric refractive index in the radio frequency region caused by convective motions. Izv. AN SSSR, Ser. geofiz. no.1:141-149 Ja '61. (MIRA 14:1)

1. Akademiya nauk SSSR, Institut prikladnoy geofiziki.
(Refraction) (Radio waves)

L 14977-63 EWT(1)/BDS AFFTC/ASE/ESD-3 RB

ACCESSION NR: AP3005438

8/0020/63/151/005/1089/1092

AUTHOR: Vul'fson, N. I.

TITLE: Influence of air humidity on the development of convection in a cloudless atmosphere

SOURCE: AN SSSR. Doklady*, v. 151, no. 5, 1963, 1089-1092

TOPIC TAGS: humidity, humidity field, atmospheric convection, humidity non-homogeneity, water-vapor, convection, thermal convection, vapor transport

ABSTRACT: Under certain conditions air humidity is more important than temperature in the development of atmospheric convection. In-flight measurements were made over the Black Sea in August 1962 with a low-inertia thermometer, a sensitive automatic dew-point hygrometer, and an accelerometer. Flights were made in the daytime in cloudless and almost windless weather. At heights of 25-50 m above the water surface, the instruments detected humidity nonhomogeneities which differed as much as 2.0-2.5 mb from the relatively uniform humidity background. Stratification of the lower layers of the air was virtually neutral. The convection phenomenon in the humidity field appears to be an important

Card 1/2

L 14977-63

ACCESSION NR: AP3005438

factor in a number of problems of atmospheric physics. For example, water vapor evaporating from the surface of the seas and oceans is transported upward primarily by convection resulting from the instability of the lower air layers. In the winter and during summer nights, when the water is warmer than the adjacent air layers, vapor is transported by thermal convection; during daytime in summer, when there is a thermally neutral or even weakly stable stratification, vapor is transported by convection associated with the humidity field. The article was presented by Academician Ye. K. Fedorov on March 15, 1963. Orig. art. has: 1 figure and 16 formulas.

ASSOCIATION: Institut prikladnoy geofiziki Akademii nauk SSSR (Institute of Applied Geophysics, Academy of Sciences SSSR)

SUBMITTED: 13Mar63

DATE ACQ: 06S:p63

ENCL: 00

SUB CODE: AS

NO REF SOV: 077

OTHER: 002

Card 2/2

VUL'FSON, N.I.; GUSAK, N.A.; SKATSKIY, V.I.

Relation of microstructure parameters to convective motions in
clouds. Izv. AN SSSR. Fiz. atm. i okeana 1 no.1:76-83 Ja '65.
(MIRA 18:5)

1. Institut prikladnoy geofiziki AN SSSR.

VUL'FSON, N.I.; IVANOV, V.N.

Structure of the temperature field in cumuli. Dokl. AN SSSR
159 no.4:786-788 D '64 (MIRA 18:1)

1. Institut prikladnoy geofiziki AN SSSR.

VULISTON, N.I.; LEVIN, L.M.

Downward convective currents. Dokl. AN SSSR 158 no.6:1329-
1333 O '64. (MIRA 17:12)

2. Institut prikladnoy geofiziki AN SSSR. Predstavleno
akademikom Ye.S. Fedorovym.

VUL'FSON, N.I.; GUTMAN, L.N.; PAVLOVA, I.S.

Effect of gravitation waves on the formation of hail clouds
in a mountainous area. Meteor. i gidrol. no.1:23-29 Ja '64.
(MIRA 17:3)

1. Institut prikladnoy geofiziki AN SSSR i Vysokogornyy
geofizicheskiy institut.

ACCESSION NR: AP4010573

S/0050/64/000/001/0023/0029

AUTHORS: Vul'fson, N. I.; Gutman, L. N.; Pavlova, I. S.

TITLE: Effects of gravitational waves on the formation of hail clouds in mountainous regions

SOURCE: Meteorologiya i gidrologiya, no. 1, 1964, 23-29

TOPIC TAGS: gravitational wave, hail cloud, cumulus cloud, wind velocity, temperature gradient, precipitation, hail

ABSTRACT: The authors' purpose is to examine the conditions under which the effects of gravitational waves may lead to the formation of vertical movements sufficiently intense to have a noticeable influence on the development of hail clouds. They consider the model of an infinitely long mountain range of arbitrary cross section and with transverse wind of constant velocity. They derive equations for air movement and compute values for different heights and breadths of the mountains. These computations show that wave forms developed by a mountainous zone may lead to the formation of strong, stationary, ascending movements of air. To test this, they investigated the relationship between development of hail in the Alazani valley and fields of temperature and wind favorable for producing atmospheric waves. For completeness and comparison they also examined temperature

Card 1/42

ACCESSION NR: AP4010573

and wind fields associated with cumulus rain clouds not producing hail as well as fields associated with cumulus clouds yielding no precipitation at all. The relationship of precipitation to wind and to temperature gradient is illustrated by Fig. 1 on the Enclosure. For winds blowing parallel to the range, regardless of temperature gradient, the weather was fair (for the four years represented by the data of Fig. 1). All kinds of clouds were observed for winds blowing at right angles to the trend of the range, but hail was more likely the nearer the wind direction was to this right-angle direction, and the higher the temperature gradient was. This means that hail is most probable under conditions most favorable for the development of gravitational waves. Orig. art. has: 3 figures and 15 formulas.

ASSOCIATION: Institut prikladnoy geofiziki (Institute of Applied Geophysics);
Vyssokogornyy geofizicheskiy institut (High-Mountain Geophysical Institute)

SUBMITTED: 00

DATE ACQ: 14Feb64

ENCL: 02

SUB CODE: AS

NO REF SOV: 002

OTHER: 002

Card 2/42

VUL'FSON, N.I.

Effect of humidity on the development of convective movements in
a cloudless atmosphere. Dokl. AN SSSR 151 no.5:1089-1092 Ag
'63. (MIRA 16:9)

1. Institut prikladnoy geofiziki AN SSSR. Predstavleno akademikom
Ye.K.Fedorovym.

(Humidity) (Atmosphere)

1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 50 51 52 53 54 55 56 57 58 59 60 61 62 63 64 65 66 67 68 69 70 71 72 73 74 75 76 77 78 79 80 81 82 83 84 85 86 87 88 89 90 91 92 93 94 95 96 97 98 99 100

101 102 103 104 105 106 107 108 109 110 111 112 113 114 115 116 117 118 119 120 121 122 123 124 125 126 127 128 129 130 131 132 133 134 135 136 137 138 139 140 141 142 143 144 145 146 147 148 149 150 151 152 153 154 155 156 157 158 159 160 161 162 163 164 165 166 167 168 169 170 171 172 173 174 175 176 177 178 179 180 181 182 183 184 185 186 187 188 189 190 191 192 193 194 195 196 197 198 199 200

201 202 203 204 205 206 207 208 209 210 211 212 213 214 215 216 217 218 219 220 221 222 223 224 225 226 227 228 229 230 231 232 233 234 235 236 237 238 239 240 241 242 243 244 245 246 247 248 249 250 251 252 253 254 255 256 257 258 259 260 261 262 263 264 265 266 267 268 269 270 271 272 273 274 275 276 277 278 279 280 281 282 283 284 285 286 287 288 289 290 291 292 293 294 295 296 297 298 299 300

301 302 303 304 305 306 307 308 309 310 311 312 313 314 315 316 317 318 319 320 321 322 323 324 325 326 327 328 329 330 331 332 333 334 335 336 337 338 339 340 341 342 343 344 345 346 347 348 349 350 351 352 353 354 355 356 357 358 359 360 361 362 363 364 365 366 367 368 369 370 371 372 373 374 375 376 377 378 379 380 381 382 383 384 385 386 387 388 389 390 391 392 393 394 395 396 397 398 399 400

401 402 403 404 405 406 407 408 409 410 411 412 413 414 415 416 417 418 419 420 421 422 423 424 425 426 427 428 429 430 431 432 433 434 435 436 437 438 439 440 441 442 443 444 445 446 447 448 449 450 451 452 453 454 455 456 457 458 459 460 461 462 463 464 465 466 467 468 469 470 471 472 473 474 475 476 477 478 479 480 481 482 483 484 485 486 487 488 489 490 491 492 493 494 495 496 497 498 499 500

501 502 503 504 505 506 507 508 509 510 511 512 513 514 515 516 517 518 519 520 521 522 523 524 525 526 527 528 529 530 531 532 533 534 535 536 537 538 539 540 541 542 543 544 545 546 547 548 549 550 551 552 553 554 555 556 557 558 559 560 561 562 563 564 565 566 567 568 569 570 571 572 573 574 575 576 577 578 579 580 581 582 583 584 585 586 587 588 589 590 591 592 593 594 595 596 597 598 599 600

601 602 603 604 605 606 607 608 609 610 611 612 613 614 615 616 617 618 619 620 621 622 623 624 625 626 627 628 629 630 631 632 633 634 635 636 637 638 639 640 641 642 643 644 645 646 647 648 649 650 651 652 653 654 655 656 657 658 659 660 661 662 663 664 665 666 667 668 669 670 671 672 673 674 675 676 677 678 679 680 681 682 683 684 685 686 687 688 689 690 691 692 693 694 695 696 697 698 699 700

701 702 703 704 705 706 707 708 709 710 711 712 713 714 715 716 717 718 719 720 721 722 723 724 725 726 727 728 729 730 731 732 733 734 735 736 737 738 739 740 741 742 743 744 745 746 747 748 749 750 751 752 753 754 755 756 757 758 759 760 761 762 763 764 765 766 767 768 769 770 771 772 773 774 775 776 777 778 779 780 781 782 783 784 785 786 787 788 789 790 791 792 793 794 795 796 797 798 799 800

801 802 803 804 805 806 807 808 809 810 811 812 813 814 815 816 817 818 819 820 821 822 823 824 825 826 827 828 829 830 831 832 833 834 835 836 837 838 839 840 841 842 843 844 845 846 847 848 849 850 851 852 853 854 855 856 857 858 859 860 861 862 863 864 865 866 867 868 869 870 871 872 873 874 875 876 877 878 879 880 881 882 883 884 885 886 887 888 889 890 891 892 893 894 895 896 897 898 899 900

901 902 903 904 905 906 907 908 909 910 911 912 913 914 915 916 917 918 919 920 921 922 923 924 925 926 927 928 929 930 931 932 933 934 935 936 937 938 939 940 941 942 943 944 945 946 947 948 949 950 951 952 953 954 955 956 957 958 959 960 961 962 963 964 965 966 967 968 969 970 971 972 973 974 975 976 977 978 979 980 981 982 983 984 985 986 987 988 989 990 991 992 993 994 995 996 997 998 999 1000

1001 1002 1003 1004 1005 1006 1007 1008 1009 1010 1011 1012 1013 1014 1015 1016 1017 1018 1019 1020 1021 1022 1023 1024 1025 1026 1027 1028 1029 1030 1031 1032 1033 1034 1035 1036 1037 1038 1039 1040 1041 1042 1043 1044 1045 1046 1047 1048 1049 1050 1051 1052 1053 1054 1055 1056 1057 1058 1059 1060 1061 1062 1063 1064 1065 1066 1067 1068 1069 1070 1071 1072 1073 1074 1075 1076 1077 1078 1079 1080 1081 1082 1083 1084 1085 1086 1087 1088 1089 1090 1091 1092 1093 1094 1095 1096 1097 1098 1099 1100

1101 1102 1103 1104 1105 1106 1107 1108 1109 1110 1111 1112 1113 1114 1115 1116 1117 1118 1119 1120 1121 1122 1123 1124 1125 1126 1127 1128 1129 1130 1131 1132 1133 1134 1135 1136 1137 1138 1139 1140 1141 1142 1143 1144 1145 1146 1147 1148 1149 1150 1151 1152 1153 1154 1155 1156 1157 1158 1159 1160 1161 1162 1163 1164 1165 1166 1167 1168 1169 1170 1171 1172 1173 1174 1175 1176 1177 1178 1179 1180 1181 1182 1183 1184 1185 1186 1187 1188 1189 1190 1191 1192 1193 1194 1195 1196 1197 1198 1199 1200

1201 1202 1203 1204 1205 1206 1207 1208 1209 1210 1211 1212 1213 1214 1215 1216 1217 1218 1219 1220 1221 1222 1223 1224 1225 1226 1227 1228 1229 1230 1231 1232 1233 1234 1235 1236 1237 1238 1239 1240 1241 1242 1243 1244 1245 1246 1247 1248 1249 1250 1251 1252 1253 1254 1255 1256 1257 1258 1259 1260 1261 1262 1263 1264 1265 1266 1267 1268 1269 1270 1271 1272 1273 1274 1275 1276 1277 1278 1279 1280 1281 1282 1283 1284 1285 1286 1287 1288 1289 1290 1291 1292 1293 1294 1295 1296 1297 1298 1299 1300

1301 1302 1303 1304 1305 1306 1307 1308 1309 1310 1311 1312 1313 1314 1315 1316 1317 1318 1319 1320 1321 1322 1323 1324 1325 1326 1327 1328 1329 1330 1331 1332 1333 1334 1335 1336 1337 1338 1339 1340 1341 1342 1343 1344 1345 1346 1347 1348 1349 1350 1351 1352 1353 1354 1355 1356 1357 1358 1359 1360 1361 1362 1363 1364 1365 1366 1367 1368 1369 1370 1371 1372 1373 1374 1375 1376 1377 1378 1379 1380 1381 1382 1383 1384 1385 1386 1387 1388 1389 1390 1391 1392 1393 1394 1395 1396 1397 1398 1399 1400

1401 1402 1403 1404 1405 1406 1407 1408 1409 1410 1411 1412 1413 1414 1415 1416 1417 1418 1419 1420 1421 1422 1423 1424 1425 1426 1427 1428 1429 1430 1431 1432 1433 1434 1435 1436 1437 1438 1439 1440 1441 1442 1443 1444 1445 1446 1447 1448 1449 1450 1451 1452 1453 1454 1455 1456 1457 1458 1459 1460 1461 1462 1463 1464 1465 1466 1467 1468 1469 1470 1471 1472 1473 1474 1475 1476 1477 1478 1479 1480 1481 1482 1483 1484 1485 1486 1487 1488 1489 1490 1491 1492 1493 1494 1495 1496 1497 1498 1499 1500

1501 1502 1503 1504 1505 1506 1507 1508 1509 1510 1511 1512 1513 1514 1515 1516 1517 1518 1519 1520 1521 1522 1523 1524 1525 1526 1527 1528 1529 1530 1531 1532 1533 1534 1535 1536 1537 1538 1539 1540 1541 1542 1543 1544 1545 1546 1547 1548 1549 1550 1551 1552 1553 1554 1555 1556 1557 1558 1559 1560 1561 1562 1563 1564 1565 1566 1567 1568 1569 1570 1571 1572 1573 1574 1575 1576 1577 1578 1579 1580 1581 1582 1583 1584 1585 1586 1587 1588 1589 1590 1591 1592 1593 1594 1595 1596 1597 1598 1599 1600

1601 1602 1603 1604 1605 1606 1607 1608 1609 1610 1611 1612 1613 1614 1615 1616 1617 1618 1619 1620 1621 1622 1623 1624 1625 1626 1627 1628 1629 1630 1631 1632 1633 1634 1635 1636 1637 1638 1639 1640 1641 1642 1643 1644 1645 1646 1647 1648 1649 1650 1651 1652 1653 1654 1655 1656 1657 1658 1659 1660 1661 1662 1663 1664 1665 1666 1667 1668 1669 1670 1671 1672 1673 1674 1675 1676 1677 1678 1679 1680 1681 1682 1683 1684 1685 1686 1687 1688 1689 1690 1691 1692 1693 1694 1695 1696 1697 1698 1699 1700

1701 1702 1703 1704 1705 1706 1707 1708 1709 1710 1711 1712 1713 1714 1715 1716 1717 1718 1719 1720 1721 1722 1723 1724 1725 1726 1727 1728 1729 1730 1731 1732 1733 1734 1735 1736 1737 1738 1739 1740 1741 1742 1743 1744 1745 1746 1747 1748 1749 1750 1751 1752 1753 1754 1755 1756 1757 1758 1759 1760 1761 1762 1763 1764 1765 1766 1767 1768 1769 1770 1771 1772 1773 1774 1775 1776 1777 1778 1779 1780 1781 1782 1783 1784 1785 1786 1787 1788 1789 1790 1791 1792 1793 1794 1795 1796 1797 1798 1799 1800

1801 1802 1803 1804 1805 1806 1807 1808 1809 1810 1811 1812 1813 1814 1815 1816 1817 1818 1819 1820 1821 1822 1823 1824 1825 1826 1827 1828 1829 1830 1831 1832 1833 1834 1835 1836 1837 1838 1839 1840 1841 1842 1843 1844 1845 1846 1847 1848 1849 1850 1851 1852 1853 1854 1855 1856 1857 1858 1859 1860 1861 1862 1863 1864 1865 1866 1867 1868 1869 1870 1871 1872 1873 1874 1875 1876 1877 1878 1879 1880 1881 1882 1883 1884 1885 1886 1887 1888 1889 1890 1891 1892 1893 1894 1895 1896 1897 1898 1899 1900

1901 1902 1903 1904 1905 1906 1907 1908 1909 1910 1911 1912 1913 1914 1915 1916 1917 1918 1919 1920 1921 1922 1923 1924 1925 1926 1927 1928 1929 1930 1931 1932 1933 1934 1935 1936 1937 1938 1939 1940 1941 1942 1943 1944 1945 1946 1947 1948 1949 1950 1951 1952 1953 1954 1955 1956 1957 1958 1959 1960 1961 1962 1963 1964 1965 1966 1967 1968 1969 1970 1971 1972 1973 1974 1975 1976 1977 1978 1979 1980 1981 1982 1983 1984 1985 1986 1987 1988 1989 1990 1991 1992 1993 1994 1995 1996 1997 1998 1999 2000

2001 2002 2003 2004 2005 2006 2007 2008 2009 2010 2011 2012 2013 2014 2015 2016 2017 2018 2019 2020 2021 2022 2023 2024 2025 2026 2027 2028 2029 2030 2031 2032 2033 2034 2035 2036 2037 2038 2039 2040 2041 2042 2043 2044 2045 2046 2047 2048 2049 2050 2051 2052 2053 2054 2055 2056 2057 2058 2059 2060 2061 2062 2063 2064 2065 2066 2067 2068 2069 2070 2071 2072 2073 2074 2075 2076 2077 2078 2079 2080 2081 2082 2083 2084 2085 2086 2087 2088 2089 2090 2091 2092 2093 2094 2095 2096 2097 2098 2099 2100

2101 2102 2103 2104 2105 2106 2107 2108 2109 2110 2111 2112 2113 2114 2115 2116 2117 2118 2119 2120 2121 2122 2123 2124 2125 2126 2127 2128 2129 2130 2131 2132 2133 2134 2135 2136 2137 2138 2139 2140 2141 2142 2143 2144 2145 2146 2147 2148 2149 2150 2151 2152 2153 2154 2155 2156 2157 2158 2159 2160 2161 2162 2163 2164 2165 2166 2167 2168 2169 2170 2171 2172 2173 2174 2175 2176 2177 2178 2179 2180 2181 2182 2183 2184 2185 2186 2187 2188 2189 2190 2191 2192 2193 2194 2195 2196 2197 2198 2199 2200

2201 2202 2203 2204 2205 2206 2207 2208 2209 2210 2211 2212 2213 2214 2215 2216 2217 2218 2219 2220 2221 2222 2223 2224 2225 2226 2227 2228 2229 2230 2231 2232 2233 2234 2235 2236 2237 2238 2239 2240 2241 2242 2243 2244 2245 2246 2247 2248 2249 2250 2251 2252 2253 2254 2255 2256 2257 2258 2259 2260 2261 2262 2263 2264 2265 2266 2267 2268 2269 2270 2271 2272 2273 2274 2275 2276 2277 2278 2279 2280 2281 2282 2283 2284 2285 2286 2287 2288 2289 2290 2291 2292 2293 2294 2295 2296 2297 2298 2299 2300

2301 2302 2303 2304 2305 2306 2307 2308 2309 2310 2311 2312 2313 2314 2315 2316 2317 2318 2319 2320 2321 2322 2323 2324 2325 2326 2327 2328 2329 2330 2331 2332 2333 2334 2335 2336 2337 2338 2339 2340 2341 2342 2343 2344 2345 2346 2347 2348 2349 2350 2351 2352 2353 2354 2355 2356 2357 2358 2359 2360 2361 2362 2363 2364 2365 2366 2367 2368 2369 2370 2371 2372 2373 2374 2375 2376 2377 2378 2379 2380 2381 2382 2383 2384 2385 2386 2387 2388 2389 2390 2391 2392 2393 2394 2395 2396 2397 2398 2399 2400

2401 2402 2403 2404 2405 2406 2407 2408 2409 2410 2411 2412 2413 2414 2415 2416 2417 2418 2419 2420 2421 2422 2423 2424 2425 2426 2427 2428 2429 2430 2431 2432 2433 2434 2435 2436 2437 2438 2439 2440 2441 2442 2443 2444 2445 2446 2447 2448 2449 2450 2451 2452 2453 2454 2455 2456 2457 2458 2459 2460 2461 2462 2463 2464 2465 2466 2467 2468 2469 2470 2471 2472 2473 2474 2475 2476 2477 2478 2479 2480 2481 2482 2483 2484 2485 2486 2487 2488 2489 2490 2491 2492 2493 2494 2495 2496 2497 2498 2499 2500

2501 2502 2503 2504 2505 2506 2507 2508 2509 2510 2511 2512 2513 2514 2515 2516 2517 2518 2519 2520 2521 2522 2523 2524 2525 2526 2527 2528 2529 2530 2531 2532 2533 2534 2535 2536 2537 2538 2539 2540 2541 2542 2543 2544 2545 2546 2547 2548 2549 2550 2551 2552 2553 2554 2555 2556 2557 2558 2559 2560 2561 2562 2563 2564 2565 2566 2567 2568 2569 2570 2571 2572 2573 2574 2575 2576 2577 2578 2579 2580 2581 2582 2583 2584 2585 2586 2587 2588 2589 2590 2591 2592 2593 2594 2595 2596 2597 2598 2599 2600

2601 2602 2603 2604 2605 2606 2607 2608 2609 2610 2611 2612 2613 2614 2615 2616 2617 2618 2619 2620 2621 2622 2623 2624 2625 2626 2627 2628 2629 2630 2631 2632 2633 2634 2635 2636 2637 2638 2639 2640 2641 2642 2643 26

10

Preparation of secondary octyl alcohol and sebacic acid.
Yu. M. Svetlov and N. S. Vul'fson. *J. Applied Chem.*
(U. S. S. R.) 9, 1613-23 (in French 1935) (1936).— A
sec. $C_8H_{17}OH$ was distd. from a mixt. of technical castor
oil (1 kg) and a soln. of 95% $NaOH$ (331 g. in 750 cc.
of water), with the reacting mixt. at 113-117° and the vapor
at 600°. One kg. of castor oil yielded 125-335 g. of crude
sec. $C_8H_{17}OH$. This product was purified by heating with
phthalic anhydride at 110-115° for 12 hrs., recrystg. the
ester (b. 55°) in gasoline (b. 60-70°), decomg. the ester
with conc. K_2CO_3 , and distg. the final product in vacuo.
Sec. $C_8H_{17}OH$, redistd. under atm. pressure, b. 178.4°, d_4^{20}
0.82008-0.82009, n_D^{20} 1.4264. The purification of the sec.
 $C_8H_{17}OH$ with 5% $KMnO_4$ in such amt. as to cause MnO_2
formation in 15-20 min. after its addn. yielded a sec.
 $C_8H_{17}OH$ b. 179.2°, d_4^{20} 0.81983-0.81985, n_D^{20} 1.4265-
1.4266. Sebacic acid was obtained as a by-product of the
above distn. After removal of the sec. $C_8H_{17}OH$, the residue
was dild. with 3 l. of hot water, half-neutralized with concd.
 H_2SO_4 (4% of the residue by vol.), and the updr. lower
layer was completely neutralized with the same acid. The
white ppt. of sebacic acid was recrystd. from 85% HCO_2H
(1:5 ratio by vol.). It m. 126-6°. Exptl. details are given.
Twenty-eight references. A. A. Podgorny

10

CA

CONDENSATION OF BROMOACETALDEHYDE WITH MALONIC ACID. N. S. Yabl'zon and M. M. Shrimyakin. *Compt. rend. acad. sci. U. R. S. S. 29, 201-7 (1940) (in English)* -- To 10 g. of malonic acid, 5 cc. of pyridine was added. Heat was evolved. When the mixt. began to cool, 12.5 g. of BrCH_2CHO was added. After 24 hrs. the reaction mixt. was heated for 2 hrs. in an oil bath at $105-15^\circ$, when CO_2 and water were given off. The mixt. was poured into 250 cc. of 5% H_2SO_4 and the flaky ppt. (about 1 g.) was filtered off and washed with ether. The product was assigned the structure of a β -keto malonic acid: $\text{BrCH}_2\text{CH}(\text{COOH})\text{COOH}$. The proof of its structure rested on the behavior of the product toward alk. hydrolyt. There should be and there was a break in the compd. and formation of BrCH_2CHO and malonic acid, which actually were recovered, the latter as Ba malonate. The product was insol. in water, alc. and benzene and very slightly sol. in ether. Rebecca Yablonsky

ASD 15.4 METALLOGICAL LITERATURE CLASSIFICATION

PROCESSING AND PROPERTY INQUIRY

BC N.S. VULFSON R-3

Fluorine of ... M. M. ... U.S.S. ...
 of ... in ... of ...
 hydrous ... of the ...
 into CO_2 , CH_2CO_2 , ...

ASB-22A METALLURGICAL LITERATURE CLASSIFICATION

SEARCH SYMBOLS

SEARCHED MAP ONLY SEE

COLLECTIONS

SEARCHED ONLY SEE

		PROCESSES AND PROPERTIES INDEX												SOL AND ATM. COEFF.											
Ca		<p>β-Lactones and β-lactonoacids. III. Chlorination of citral with melenic acid. N. B. Vulfson and M. M. Shemyakin, <i>J. Gen. Chem.</i> (U.S.S.R.), 13, 436-47 (1943) (English summary); cf. C. A. 37, 611P. —Melenic acid (20 g.), 30 g. citral and 3 cc. glacial AcOH concentrated with 3 cc. piperidine, heated on a water bath for 1.5 hrs., when the mass solidifies. After washing with Et₂O, the solid is crystd. from 60% alc., to yield 10 g. (45%) of 2,2-dilactone of <i>cis</i>-pyridenemalenic acid, Me₂C·CH(CN)₂·CO·O—CO</p> <p>Me₂C·CH₂·CH·CH·CO·O (I), m. 180-7°. The mother liquor yields <i>cis</i>-pyridenemalonic acid, bp. 165-75° (at 0.5 mm.), 1 (1 g.), heated to soln. with 4 cc. 10 N NaOH, boiled for 2-3 min. and acidified by dil. H₂SO₄, yields 0.55 g. MeC·CH(CH₂·CH₂·CM₂(OH)CH₂·CH·CH(CO)H·CO₂H, m. 113-14° (after extra. with hot benzene); the crvstl. benzene extract yields 0.1 g. Me₂C·CH(CH₂·CH₂·CM₂(OH)CH₂·CH·CH₂·CO₂H (II), m. 120-0.5°, a soln. of the Na salt of the above acid, on acclification, yields the mixt. of the acid and the above lactone, indicating fairly ready decarboxylation; the Ag salt of the acid appears to have an appreciable water soly., while AcCl leads to loss of some CH₂ and yields I. I boiled with 10 N NaOH and acclified, yields 67% II, which, treated with slight excess of 0.1 N NaOH, acclified by HNO₃ and treated with AgNO₃, gave MeC·CH(CH₂·CH₂·CM₂(OH)CH₂·CH·CH(CO)H₂. II is not affected by Ar₂O in pyridine at about 100°, while AcCl yields 4-5% MeC·CH(CH₂·CH₂·CM₂(OAc)CH·CH·CH₂·CO₂H in 117-18° (from benzene). Quilization of II by alk. KMnO₄ yielded (CO₂H)I, while no HCO₂H was detected, thus indicating the presence of the β-lactone ring. IV. The rate of opening of the β-lactone ring. M. M. Shemyakin and N. B. Vulfson, <i>Ibid.</i>, 448-56 (English summary). —The authors studied the rate of ring opening of the lactones Me₂C·CH(CO₂H)·CO₂H and PhCH·CH(CO₂H)·CO₂H at various temps. The results, presented in tabular and graphical form, show that the ring opening of β-lactones is a first-order reaction. There are indications that the hydrolysis in alk. medium leads to unsat. acids and that the hydrolysis reaction is a tautomeric one. For the 2 lactones tried (above) the activation energy of hydrolysis was found to be 19,323 and 22,532 cal., resp. O. M. Koonin.</p>																							
		<p>A-B-S-A METALLURGICAL LITERATURE CLASSIFICATION</p> <p>SOURCE SYLLABUS</p> <p>SYNOPSIS AND ONLY USE</p> <p>REVISIONS</p> <p>DATE REVISOR</p> <p>REVISOR'S NAME AND TITLE</p>																							

VULFSON, N. S.

"Investigation of B-Lactones and B-Lactono-Acids. IV. Study of the Reaction Rate of the Cleavage of the Lactone Ring". Shemyakin, M. H. and Vulfson, N. S. (p. 455)

SO: Journal of General Chemistry (Zhurnal Obshchei Khimii) 1943, Volume 13, no. 6.

PRECEDENCE AND PRIORITY INDEX																	
<p><i>ca</i></p> <p>Chemical nature of vitamin A. N. S. Val'son. <i>Us-</i> <i>spek Khim.</i> 13, 437-61(1944); cf. C.A. 37, 6657.-A review of the chemistry of vitamin A and vitamin A, and the carotenes, including degradation, synthesis, and analytical methods, with 130 references. S. Gottlieb</p>																	
<p>A.S.S.L.A. METALLURGICAL LITERATURE CLASSIFICATION</p>																	
<p>FROM SOURCE</p>																	
<p>COLLECTION</p>																	
<p>DATE OF RECEIPT</p>																	

CA

Research in β -lactones and β -lactono acids. V.
Mixed anhydrides of malonic and acetic acids. N. S.
Vul'fon, *J. Gen. Chem.* (U.S.S.R.) 19, No. 10, 2309-
81 (1949) (English translation).—See C..I. 44, 1901f.
E. J. C.

CA

β Lactones and *α* lactono acids. V. Mixed anhydrides of malonic and acetic acids. N. E. Vol'kov. *Zhur Obshch. Khim.* (J. Gen. Chem.) 19, 1001 (1940); cf. C.A. 34, 3235. $\text{CH}_3(\text{CO}_2\text{H})_2$ with Ac_2O yields 2 mixed *aceto-malonic anhydrides* in a reaction catalyzed by H_2SO_4 . Shaking 10 g. $\text{CH}_3(\text{CO}_2\text{H})_2$, 40 g. Ac_2O , and 3 drops H_2SO_4 until solid occurs and letting stand overnight, followed by concn. *in vacuo* at 40–50°, gave a sirup which on treatment with 15 ml. abs. EtOH, followed by cooling and extr. with Et₂O, gave 1.5 g. MeOAc and 10%, $\text{CH}_3(\text{CO}_2\text{Me})_2$ (I), b. 74–75°, n_D²⁰ 1.4140, while an ext. with Na₂CO₃ gave 21.7% $\text{HO}_2\text{CCH}_2\text{CO}_2\text{Me}$ (II), b. 115–8° (decompn.). If the reaction mixt. above after vacuum concn. is extr. with Et₂O and the ext. is treated with MeOH, there is formed 2.6 g. MeOAc, 81% I, and 22% II, as well as a trace of $\text{CH}_3(\text{CO}_2\text{H})_2$. Similar results are obtained if H_2SO_4 is omitted and the mixt. is allowed to stand 24 hrs. before concn. and reaction with MeOH. $\text{CH}_3(\text{CO}_2\text{H})_2$ (12.8 g.) with 0.3 g. AcCl in Et₂O gave upon filtration a yellow sirup, which gave 91.5% I with MeOH; 8.4 g. Ag salt and 3.1 g. AcCl gave 84.7% II and 11.5% MeOAc. The mixed anhydride from 10 g. $\text{CH}_3(\text{CO}_2\text{H})_2$ and 10 g. Ac_2O gave with 20 g. BuOH, 2.7 g. $\text{CH}_3(\text{CO}_2\text{H})_2$, 39% $\text{HO}_2\text{CCH}_2\text{CO}_2\text{Bu}$ (undistillable without decompn.), and 11.5% di Bu ester, as well as 3.5 g. BuOAc; similar reaction with 12 g. Me₂COH gave 35.7% $\text{HO}_2\text{CCH}_2\text{CO}_2\text{Me}$ (isolated as the Ag salt), and 14.5% di-*tert*-Bu ester, b. 230–5° (with some decompn.), as well as 1 g. *tert*-BuOAc; 2-octanol (25 g.) gave 2.2 g. $\text{CH}_3(\text{CO}_2\text{H})_2$, 0.5 g. 2-octyl acetate, 14.5% 2-octyl II malonate, and 15.9% corresponding neutral ester, b. 100–70°, n_D²⁰ 1.4367. Dodecylalc. (35 g.) gave a small amt. of dodecyl acetate, b. 140–52°, 16.5% didodecyl malonate, m. 33–4°, and 23% dodecyl II malonate, m. 42–3° (from iso-Am₂O); 20 g. PhOH gave a little PhOAc, 13.2% di-Ph malonate, m. 49.5–51.0°, and 17.3%

Ph II malonate, m. 65–67° (from iso-Am₂O), while 10 g. PhNH₂ gave 71% malonanilide, m. 223–4.5° (from MeOH); similar addn. of 15 g. PhNH₂ in 25 ml. Et₂O gave 12 g. of the anilide while the Et₂O mother liquor yielded about 1 g. AcNHPh and the alk. ext. gave 4.5 g. malonanilide, m. 140–2° (from AcOH), which on heating to the mp. gave AcNHPh. Addn. of 2 drops H_2SO_4 to 5 g. II and 20 ml. Ac₂O, letting stand 24 hrs., and evapn. *in vacuo* at 50° gave the mixed anhydride of acetic acid and II, b. 70–1°, n_D²⁰ 1.4100, which (3 g.) treated with 5 ml. MeOH gave I, while 3 g. PhNH₂ gave 0.4 g. AcNHPh and 1 g. II. Treatment of the mixed anhydride with p-CH₃NC₆H₄CH₃ in hot up. alc. NaOH for 1 hr. gave p-nitrobenzyl acetate and malonate, m. 77–9° and 82–5–84.0°, resp. Allowing 1 g. $\text{HO}_2\text{CCH}_2\text{CO}_2\text{Bu}$ and 16 g. Ac₂O to stand 24 hrs. gave 30% mixed anhydride of acetic acid and Bu II malonate, b. 116–18°, which (1 g.) with 2 g. PhNH₂ gave AcNHPh. Similarly $\text{HO}_2\text{CCH}_2\text{CO}_2\text{Ph}$ gave the corresponding mixed anhydride with 1 g. II, m. 55.5–56.0° (from iso-Am₂O), giving $\text{HO}_2\text{CCH}_2\text{CO}_2\text{Ph}$ and AcNHPh with PhNH₂. G. M. K.

YDD PA 169T28

VULFSON, N. S.

Sep 50

USSR/Chemistry - Laboratory Equipment

Continuous-Action Counterflow Extraction Apparatus for Laboratory Use,"
F. N. Stepanov, N. S. Vulfson, I. A. Mikova, Sci Res Inst of Org Intermediate
Products and Dyestuffs

"Zavod Lab" Vol XVI, No 9, pp 1131.

New apparatus for extraction from solutions was constructed and tested in operation. Extraction occurs in narrow vertical tube in which solution and solvent, flowing toward each other, are stirred vigorously by spiral ribbon mixer. Apparatus is designed for operation with types of solvent heavier or lighter than solution. Main advantage of extractor is its efficiency, and small amount of solvent used.

PA 169T28.

VUL'FSON, N.S.; PODREZOVA, T.N.; SENYAVINA, L.B.

Dieckmann reaction. Part 13: Infrared and ultraviolet spectra of methyl- and carbethoxy derivatives of 3-chromanone. Zhur. ob. khim. 34 no.8:2676-2681 Ag '64. (MIRA 17:9)

1. Institut khimii prirodnikh soyedineniy AN SSSR i Nauchno-issledovatel'skiy institut organicheskikh poluproduktov i krasiteley (NIOPiK).

PUCHKOV, V.A.; STEPANOV, V.M.; VUL'FSON, N.S.; ZYAKUN, A.M.; KRIVTSOV, V.F.

Mass spectrometry of amino acid methylthiophydantoins. Dokl.
AN SSSR 157 no.5:1160-1163 Ag '64. (MIRA 17:9)

1. Institut khimii prirodnikh soyedineniy AN SSSR.

C.A.

10

β-Lactones and *β*-lactono acids. VI. Mechanism of formation of *β*-lactono acids. N. S. And'ronov, *Zhur. Obshch. Khim.* (J. Gen. Chem.) 20, 1233 (1950); cf. C. 44, 6192a. It has been shown that only the neutral malonic acetic anhydride (I) is capable of reacting with carbonyl derivs.; the mixed acetic anhydride does not react. H₂SO₄, which catalyzes the anhydride formation, does not participate in the reaction with CO derivs. The 1st step of the reaction of I with Me₂CO is the formation of the mixed anhydride of AcOH and Me₂C(OH)CH₂(CO₂H), which undergoes an intramol. reaction yielding the lactono-acid and Ac₂O. Crude I and Me₂CO, allowed to stand overnight, readily yield 18.6-33.5% isopropylidene-malono-*β*-lactone, RR'C(CH₂CO₂H)CO₂(R, R' = Me)

m. 96.7° (from Me₂CO or C₆H₆), also obtained in 48.6% yield from 6.4 g. CH₃(CO₂Ag), in 10 g. dry Me₂CO with 10 g. AcCl (added dropwise), followed by filtration and standing overnight; BzCl instead AcCl gives the same product, in addn. to some BzOH (amts. unstated). The crude I from 10 g. CH₃(CO₂H), and 10 g. BzH, let stand overnight, gave 27.0%; benzylidenemalono-*β*-lactone, m. 145.6° (decolor); from Me₂CO-C₆H₆, also obtained (1 g.) by addn. of 6.4 g. CH₃(CO₂Ag) to 10 g. BzH, followed by 5 g. AcCl. m-ONC₆H₄CHO in the 1st reaction gave 0.4 g. m-NO₂ analog, m. 154.5-59.0° (from MeOH), while cyclohexanone (10 g.) gave 1.5 g. cyclohexylidenemalono-*β*-lactone, m. 84.5°. The lactono acids were isolated in the form of the resp. Ag salts (undescribed and used only for analyses). When Me₂C(CH₂CO₂H)₂ was treated with a trace of H₂SO₄ in Ac₂O, no lactonization took place even in 3 days, nor did its di-Ag salt yield any lactone with AcCl in Me₂CO; the benzylidene analog behaved similarly.

Addn. of 3 drops concd. H₂SO₄ to 6.2 g. Me₂C(CH₂CO₂H)₂ in 25 ml. Ac₂O, followed by 2 hrs. at 60° and standing for 2 days gave, after distn. of the Ac₂O in vacuo and washing the residue with Na₂CO₃ soln. (in Et₂O), 4 g. isopropylidene-acetic anhydride, b_p 140-2°, b_p 117-18°, which yields the anilide, m. 127.5-8.0° (from EtOH); 0.5 g. original acid is reclaimed. Me₂C(CH₂CO₂Ag) with AcCl in Et₂O gave only the free acid, m. 67.5-9.0°. Addn. of 10 g. AcCl to 6.4 g. CH₃(CO₂Ag) in 15 ml. dry Me₂CO, followed by filtration and sepn. of the filtrate into parts (a) and (b) gave: from part (a), allowed to stand 2 hrs. after filtration, an unstated amt. of CH₃(CO₂H), and from part (b), allowed to stand 1 day, an unstated amt. of isopropylidenemalono-*β*-lactone. A similar reaction in which the 24-hr. filtrate was treated with dry MeOH gave MeCOAc, CH₃(CO₂Me), AcOH, and a small amt. of the above lactone. PhOH instead of MeOH gave di-Ph malonate, m. 185.0-5°. Distn. of the 24-hr. filtrate yielded a small amt. of Ac₂O and the above lactone. VII. Relationship between structure of carbonyl compounds and the product of its condensation with malonic acid. *Ibid.* 445-44. The reactions of neutral acetic malonic anhydride (I) with CO derivs. indicate that the presence of an actual or potential OH group in the carbonyl compd. leads to formation of malonic esters and not the lactono acids; the latter form predominantly if the carbonyl deriv. is capable of resonance, with considerable local resonance in the CO group. The condensation does not occur if many resonant structures are possible in which the pos. pole is not at the C atom. I [from 10 g. CH₃(CO₂H)] and 10 g. AcPh in 48 hrs. at room temp. gave 2.5 g. 2-phenylthylidenemalono-*β*-lactone, m. 91.5-2.0° (Ag salt of free acid used in analysis). Menthone

(over)

(15 g.) in a 24-hr. reaction gave 4.8 g. *menthylidenemalonate*- β -lactone, needles, m. 107.6-8.0° (decomp.; from MeOH); both the Ag salt of the lactone and di-Ag salt of the free acid were analyzed. PhCH:CH:CHO (13.5 g.) gave in 24 hrs. 3.5 g. *hydrocinnamylidenemalonate*- β -lactone, m. 130-40° (from MeCO-MeOH) (Ag salt of the lactone). *o*-Cl-C₆H₄CHO (7 g.) gave 1.5 g. *o*-chlorobenzylidenemalonate- β -lactone, m. 1210-135° (from CHCl₃) (the Ag salt); *p*-Cl-lactone, m. 1210-135° (from CHCl₃) (the Ag salt); *m*-Cl-analog (1.1 g.), obtained similarly, m. 151-5° (from EtOH and Me₂CO) (Ag salt); *m*-Cl-analog, m. 161-2° (from EtOH and C₆H₆), was obtained in 2.5-g. yield (Ag salt). Diacetylene alc. (20 g.) yielded 1 g. acid ester of diacetylene alc. and CH₃(CO₂H)₂. AcCH₂CM₂OCOCCH₂CO₂H, an odoriferous yellow oil, yielding a semicarbazone, m. 232-4° (from EtOH). Paraldehyde (15 g.) gave 4.5 g. acetaldoxime malonate, C₆H₁₀O₄, an oil yielding a semicarbazone, m. 152-3° (from EtOH); AcH reacts similarly, and the same product forms from AcCl, AcH, and CH₃(CO₂Ag). EtCHO (20 g.) gave 5 g. orange propionaldoxime malonate, C₆H₁₀O₄; semicarbazone, m. 177-0° (from EtOH). I [from 30 g. CH₃(CO₂H)] and 45 g. AcCH₂CH₂CH₂ gave 15 g. of the corresponding enol acid malonate, C₆H₁₀O₄, a yellow oil, yielding the *p*-nitrobenzyl ester, m. 212.5-4.0°. I [from 10 g. CH₃(CO₂H)] and 15 g. mesityl oxide gave 3 g. of enol acid malonate, oil, C₆H₁₀O₄ (Ag salt), also obtained from AcCl, CH₃(CO₂Ag), and mesityl oxide. Pseudoionone (10 g.) with I gave 3 g. of the enol acid malonate, oil, C₆H₁₀O₄; Ag salt. PhCH:CHCOMe (15 g.) gave the enol acid malonate (4 g.), oil (Ag salt). I and 15 g. Cl₂CCO gave in 24 hrs. 8 g. chloral diacrylate, t, 111.5-12.5°, b, 98-9°.

d₄²⁰ 1.4214; reaction of 6.4 g. CH₃(CO₂Ag)₂ with 6 g. Cl₂-CCO and 10 g. AcCl gave in 24 hrs. 3 g. chloral chloroacetate, ClCCOAc, b. 181-7°, d₄²⁰ 1.4773. Chloral gave mostly a tar and a little *p*-cymene; furfural gave a tar, while PhCHO and *p*-MeC₆H₄CHO failed to react. VIII. Condensation of cinnamaldehyde with malonic acid. *Ibid.* 505-9.—Crude acetic malonic anhydride (from 10 g. malonic acid) and 12.2 g. PhCH:CHCHO, let stand 1 day, yielded 6 g. malonic acid and 3 g. *cinnamylidenemalonate* acid (I), m. 205-6°, sepl. by CHCl₃ extn.; evapn. of the mother liquor yielded 3 g. PhCH:CHCH(CO)₂C:CHCH:CHPh, (II), m. 154-1.5° (from ligroin-C₆H₆), also obtained in 7.5-g. yield (m. 155.3-5.7°) from 3 drops concd. H₂SO₄, 10.4 g. malonic acid, 25 g. Ac₂O, and 14 g. PhCH:CHCHO after 1 day's standing; some I also forms. Titration of II uses 1 equiv. of NaOH and evapn. of the C₆H₆ ext. of the reaction soln. regenerates II; if the product, however, is boiled with alkali, 2 equivs. of NaOH are used and acclification yields I. Pure I, m. 156.5-7.5°, is obtained in 0.7-g. yield by treating 0.6 g. PhCH:CHCHO, 2 g. AcOH, and 2 g. Ac₂O with 0.6 g. I and 1 drop of concd. H₂SO₄ and letting stand 1 day. AcCl (5 g.), 0.6 g. CH₃(CO₂Ag)₂, and 2.7 g. PhCH:CHCHO let stand 1 day and the AgCl removed gave 1.4 g. *cinnamylidenemalonate*- β -lactone, m. 132-3° (from Me₂CO); addn. of AgNO₃ to the neutralized soln. gave the Ag salt, C₆H₁₀O₄Ag; the salt formed after boiling the acid in alk. soln. was C₆H₁₀O₄Ag. Boiling the lactone with Me₂CO 2 hrs. gave none I (from MeOH) and unchanged lactone. Cinnamaldehyde diacrylate, m. 84-5°, prepl. in 11.5-g. yield from 10 g. PhCH:CHCHO and 20 g. Ac₂O with a trace of H₂SO₄ is unstable on storage. IX. Decarbonyla-

C. A.

(cont'd)

tion of β -lactono acids. *Ibid.* 603-2. Heating 1 g. isopropylidenemalonate- β -lactone with 10 ml. Ac_2O 25 hrs. to $60-70^\circ$ gave on distn. in *vacuo*, 0.3 g. β , β -dimethylacrylic acid, m. $64-6^\circ$, also obtained (0.2 g.) on similar treatment of 1.5 g. $\text{Me}_2\text{C}:\text{C}(\text{CO}_2\text{H})_2$. Heating 3.5 g. benzylidenemalonate- β -lactone with 35 ml. Ac_2O 20 hrs. at $60-70^\circ$ similarly gave 1.5 g. cinnamic acid. The formation of unsatd. acids from thermal treatment of the β -lactono acids in Ac_2O is an apparently general reaction and Minot's [*Ber.* 27, 1225(1904)] synthesis of $\text{Me}_2\text{C}:\text{CHCO}_2\text{H}$ from Me_2CO and malonic acid with Ac_2O probably involves β -lactono acid formation. X. β -Lactone of menthylideneacetic acid. *Ibid.* 603-8. Dehydration of Et menthylacetate according to Wallach (*Ann.* 323, 151(1902); *C.A.* 1, 2100) gave Et menthylacetate, b. 125.7° , and menthylideneacetic acid - β -lactone, $\text{H}_2\text{C}:\text{CH}(\text{Me})\text{CH}_2\text{CH}_2\text{CH}(\text{iso-Pr})\text{CO}_2\text{O}$, b. $138-41^\circ$; the latter opens the ring on

heating with 0.1 N NaOH and acidification of the mixt. yields a product which is again not titrated by cold NaOH; i.e. the ring opening is reversible. A soln. of AgNO_3 to carefully neutralized (HNO_3) soln. which had been boiled 0.5 hr. with excess NaOH gave the Ag salt of the acid, $\text{C}_{11}\text{H}_{17}\text{O}_4\text{Ag}$. Oxidation of the lactone with KMnO_4 in Me_2CO gave menthone, the same being obtained from Et menthylacetate; menthone is not affected. When the lactone is boiled with 60% KOH, menthone is formed. The results indicate that Wallach's proposed structure of the lactone (γ -lactone) is incorrect and that β -lactone is the correct formulation.

SSR/Chemistry-Beta -Hydroxy Acids

Apr 52

"The Problem of Preparing β -Hydroxyacids; Some Remarks on the Articles of A. M. Gakhokidze "Condensation of Ketones With Esters of Organic Acids," N. S. Vul'fson

"Zhur Obshch Khim" Vol XXII, No 4, pp 718-720

In repeating Gakhokidze's experience producing β -hydroxy acids, his findings were not confirmed. The constants of the original malonic ester cited by Gakhokidze do not agree with the constants of the diethyl acid dimethyl ester of isopropylidene malonic acid cited by the authors do not correspond to the

224154

constants given for this comp. Gakhokidze and Vul'fson's suggested method for separating hydroxy acids by saponification with alcoholic alkali was not applied, because these substances decompose easily.

VUL'FSON, N.S.

224154

VUL'FSON, N. S.

Defended his Dissertation for Doctor of Chemical Sciences, Kazan' State University
1953

Dissertation: "Investigation of Beta-Hydroxy Derivatives of Malonic Acid"

SO: Referativnyy Zhurnal Khimiya, No. 1, Oct. 1953 (W/29955, 26 Apr 54)

~~SECRET~~
VUL'FSON, N. S.

The Reformatsky reaction with bromomalonate ester.
Condensation of benzaldehyde with diethyl malonate.
N. S. Vul'fson, *Sbornik State Obshchestvennoy Akad. Nauk S.S.S.R.* 1, 523-6 (1953). To 115 g. dry MgCO_3 and 160 g. $\text{CH}_2(\text{CO}_2\text{Et})_2$ in 200 ml. abs. Et_2O was added at 0° 20 g. powd. KOH (temp. rise to $8-12^\circ$) and the mixt. kept 2 hrs. in the cold, then 10 days at room temp. with occasional shaking, gave 100 g. K Et malonate. No condensation was achieved, since the liquid portion of the mixt. consisted of $\text{CH}_2(\text{CO}_2\text{Et})_2$ only (cf. Galokidze and Guntsadze, C.A. 42, 2570i and V., C.A. 47, 1704h). Refluxing 10.5 g. NaI , 24 g. $\text{BrCH}_2\text{CO}_2\text{Et}$, and 1. dry CaH_2 and 6.55 g. Zn shavings (pretreated with 10% HCl , washed and heated with a little iodine) 3 hrs. gave after cooling and acidification with dil. H_2SO_4 an orange oil, which suffers decompn. during attempted distn. *in vacuo* yielding a distillate which is brightly colored (red or blue); re-distn. gave 10 g. dl-Et benzalmonate, bp $185-7^\circ$, readily hydrogenated to the dihydro deriv., which is hydrolyzed to the free acid, m. $118-17^\circ$. Free benzalmononic acid m. $191-2^\circ$. If the mixt. (above) is saponified directly with 5% NaOH in the cold in MeOH , there is obtained a solid ppt. which treated with dil. H_2SO_4 also gave benzalmononic acid. The product which had not been heated (e.g., in the course of distn., above) does not take up H_2 . Thus the directly formed intermediate must be a β -hydroxy deriv., which loses H_2O on heating and yields the benzalmonate. The HIO deriv. could not be isolated in the pure state. Et H benzalmonate, m. $91-2^\circ$, is formed in small amounts during the above reaction.
G. M. Kosolupoff

Handwritten signature

VOL. FSON - N.S.

β -lactones and β -lactono acids. XI. Reformatskii
reaction with bromomalonic ester. 2. β -Lactone of
benzylmalonic acid. N. S. Vulf'son. J. Gen. Chem.
U.S.S.R. 24, 1917-18(1054) (for translation).—See C.A.
49, 13108c. B. M. R.

Vul'fson, N. S.

USSR/Chemistry - Reaction processes

Card 1/1 Pub. 151 - 28/37

Authors : Vul'fson, N. S.


Title : Study of beta-lactones and beta-lactonic acids. Part 11.- The Reformatzky reaction with bromomalonic ester. II. Beta-lactone of benzalmalonic acid


Periodical : Zhur. ob. khim. 24/10, 1853-1855, Oct 1954

Abstract : The derivation of beta-lactone of benzalmalonic acid through the reaction of acetyl chloride with sodium salts formed during the saponification of the benzaldehyde - bromomalonic ester condensation (according to Reformatzky), is described. The product derived from a Reformatzky reaction between benzaldehyde and bromomalonic ester, and the intermediate product formed during the formation of beta-lactonic acid, are listed. Four references: 3-USSR and 1-USA (1936-1953).

Institution : The K. E. Voroshilov Scientific Research Institute of Semi-Products and Dyes

Submitted : May 8, 1954





USSR.

Synthesis of exobutenol. N. S. Voznesenskiy (K. E. Voznesenskiy), Inst. Org. Intermed. and Catalysis, Akad. Sci. USSR, Leningrad. Prikl. Khim., 21, 1111-1112 (1951). 1400 g. Me₂CO and 370 g. 37% formalin was added gradually about 6 hr. 4% NaOH until pH 7.5 was reached; over 1 hr., the soln. was kept, with appearance of yellow color and a slight rise in temp., more NaOH being added if needed to maintain alkalinity; after 20 hr. at 33°, 6 g. hydroquinone was added, stirred briefly, and heated rapidly to reflux. Me₂CO distn. to 82-83° in the reaction flask, followed by filtration and distn. of the liquid; gave after 2 fractional distns. 66-68% 4-butenol-2 and 3-6% 1-butanol-2. The best results are obtained at pH 7.5 for 3-6 hrs. total duration. Reduction or elimination of the amt. of hydroquinone listed above gave lower yields in the right. G. M. Kozlov.

Vul'fson, N. S.

6


✓ Preparation of 4-nitrosalicylic acid. V. N. Sitnikov, N. S. Vul'fson, and I. A. Shkova (K. B. Voshchikov Sci. Res. Inst. Org. Synth. and Dyes, Moscow). *Chem. Abstr.* 57, 13433 (1954). — To 100 ml. 25% NaOH, 500 ml. H₂O, and 150 g. ice was added in 1 hr. 50 g. 4-nitrophthalimide, stirred 1 hr. at 5-8°, treated over 20 min. with 120 ml. NaOCl soln. (14% active Cl), stirred 1 hr., heated to 60°, allowed to cool slowly over 18 hrs., filtered and acidified, yielding 80% mixed nitroanthranilic acids, m. 237-41°. This (50.5 g.) added at 97° to 300 ml. 50% H₂SO₄, cooled to 3°, diazotized with 57 ml. 40% NaNO₂, stirred 0.5 hr. and added to 350 ml. H₂O and 470 g. CuSO₄, kept at 116-8°, stirred 1 hr., filtered hot and washed with H₂O, gave 87.1% mixed nitrosalicylic acids, m. 183-204°. This (35 g.) in warm soln. of 10 g. NaHCO₃ in 500 ml. H₂O, was filtered, heated to 97-5°, acidified with 18 ml. 18% HCl and cooled, yielding 18 g. crude product, which crystd. from 300 ml. 30% EtOH gave 16 g. pure 4-nitrosalicylic acid, m. 219-30°. O. L. Kosolapoff

(2)

gaw

HENRY, Thomas Anderson; DITKOVSKIY, D.P. [translator]; SUVOROV, M.N., [translator]; RODIONOV, V.M., akademik, redaktor [deceased]; VUL'FSON, N.S., doktor khimicheskikh nauk, redaktor; LEVINA, M.M., otvetstvennyy redaktor; SHEPAK, Ye.G., tekhnicheskii redaktor

[The plant alkaloids. Translated from the English] Khimiia rastitel'nykh alkaloidov. Perevod s angliiskogo. Pod red. V.M. Rodionova. i N.S.Vul'fsona. Moskva, GIs. nauchno-tekhn. izd-vo, khim. lit-ry, 1956. 904 p. (MLRA 10:1)
(Alkaloids)



Vul'fson, N.S.

Chem

✓ Co-condensation of benzaldehyde with bromoacetonitrile.
N. S. Vul'fson and L. Kh. Ginosrad. *Dokl. Akad. Nauk SSSR* 106, 569-70 (1966). — To a refluxing soln. of 21.3 g. BaH_2 , 32 ml. C_6H_6 , and 8 ml. Et_2O containing 18 g. activated Zn dust there was added in 0.5 hr. 24 g. BrCH_2CN . After refluxing 1 hrs. the cooled mixt. was stirred 1 hr. with 120 ml. 10% H_2SO_4 , filtered, and the org. layer extd. with 10% H_2SO_4 and H_2O , and the combined layers extd. with C_6H_6 and combined with the original org. layer. Distn. gave 14.2 g. $\text{PhCH(OH)CH}_2\text{CN}$, b.p. 117-61°. This shaken with 12% H_2O in the presence of 1 g. NaOH and little EtOH gave the corresponding amide, m.p. 120° (from EtOH). Refluxing the nitrile with 4 g. NaOH gave 63.8% $\text{PhCH=CHCO}_2\text{H}$.
G. M. Fozolapoff

Chem

Vul'fson, N.S.

Distr. 4E41

✓ Dieckmann reaction. I. Synthesis of α -phenyleneacetic acid- β -propionate and its ester, labeled with carbon-14 in the carboxyl group of the propionic acid. N. S. Vul'fson and M. O. Iodko. *Ukrain. Khim. Zhur.* 23, 408-41 (1967) (in Russian). 2. $\text{EtC}_6\text{H}_4\text{COOH}$ and $\text{NaC}_6\text{H}_4\text{CO}_2\text{Et}$ form Ia (CH_3OH) contg. CuCN and Cu at $140-60^\circ$ 3,2-HOC $\text{CH}_2\text{CH}_2\text{CH}_2\text{CO}_2\text{Et}$ lyzed to 5,2-HOC $\text{CH}_2\text{CH}_2\text{CH}_2\text{CO}_2\text{H}$, m. $216-18^\circ$, converted by Na and AcOH in α -EtO $\text{CCH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CO}_2\text{Et}$ (I), b.p. $191-3^\circ$. II. Mechanism of the cyclization of diethyl α -phenyleneacetic- β -propionate. P. N. Sigmoy, M. O. Iodko, and N. S. Vul'fson. *Ibid.* 23, 412-42. Cyclization of I with Na in C_6H_6 contg. EtOH form α -1,2,3,4-tetrahydro-3-oxo-2-naphthoate-carboxy- C^{14} (II), b.p. $103-4.5^\circ$, n_D^{20} 1.5198; Cu deriv., m. $164-5^\circ$. When hydrolyzed and decarboxylated 3,4-dihydro-2(1H)-naphthalenone-2- C^{14} , t. $83-5^\circ$, was formed. This shows that the ring formation is between the CO_2Et group of the propionic acid and the H of the acetate. The Na deriv. of II and MeI form Et 1,2,3,4-tetrahydro-1-methyl-2-oxo-1-naphthate, b.p. $113-15^\circ$, n_D^{20} 1.5140, hydrolyzed and decarboxylated to 3,4-dihydro-1-methyl-2(1H)-naphthalenone, m. $107-12^\circ$, n_D^{20} 1.5400, semicarbazone, m. $193-6^\circ$.
John H. McCarty //

017

Vul'fson, N.S.

STEPANOV, F.N.; IODKO, M.O.; VUL'FSON, N.S.

Investigation of Dieckmann's reaction. Part 2: Mechanism of the
cyclization reaction of diethyl ether of o-phenylacetic- β -propionic
acid. Ukr.khim.zhur. 23 no.4:489-492 '57. (MIRA 10:10)
(Chemical reaction--Mechanism)
(Propionic acid)

VUL'FSON, N.S.

79-2-21/64

AUTHORS: Vul'fson, N. S. , Zaretskiy, V. I.

TITLE: Investigations in the Field of Dieckmann (Dikman)'s Reaction
(Issledovaniye v oblasti reaktsii Dikmana)
III. Cyclization of the Diethyl Ether of α -Methylpinelic Acid
(III. Tsiklizatsiya dietilovogo efira α -metilpinelinyoy kisloty)

PERIODICAL: Zhurnal Obshchey Khimii, 1958, Vol. 28, Nr 2, pp. 369 - 372 (USSR)

ABSTRACT: The cyclization of the diethyl ether of α -methylpinelic acid was described by Dieckmann (Dikman) (reference 1). He supposed that 2-methyl-6-carbethoxycyclohexanone is formed on that occasion, but it was not eliminated in a pure state. As a continuation of the investigation of the cyclization process of the unsymmetrical dicarboxylic acid ether (reference 2), according to Dikman, the authors wanted to determine the structure of the β -keto ether. The latter is produced by the cyclization of the diethylether of α -methylpinelic acid. The keto ether (I) is obtained as the only product. The formation of β -keto ether could not be determined. It is assumed that the exclusive formation of β -keto ether (I) in the cyclization is a consequence of the increased electron density at the α -carbon atom under the influence of the I-effect of the CH_3 group. This renders the dissociation of the proton and the formation of

Card 1/2

79-2-21/64

Investigations in the Field of Dieckmann (Dikman)'s Reaction . III. Cyclization
of the Diethyl Ether of α -Methylpimelic Acid

the organic anion, which are necessary for the cyclization, difficult. Summary: The authors investigated the cyclization of the diethyl ether of α -methylpimelic acid according to Dikman. It was shown that 2-methyl-6-carbethoxycyclohexanone is the primary reaction product; the formation of 2-methyl-6-carbethoxycyclohexanone was not observed. The structure of 2-methyl-6-carbethoxycyclohexanone was confirmed by the production of 2,6-dimethylcyclohexanone which was obtained by methylation with methyl iodide and subsequent hydrolysis and decarboxylation. There are 13 references, 2 of which are Slavic.

ASSOCIATION: Scientific Research Institute for Organic Semiconductors and Dyes
(Nauchno-issledovatel'skiy institut organicheskikh poluprovodnikov i krasiteley)

SUBMITTED: January 21, 1957

AVAILABLE: Library of Congress

Card 2/2

VUL'FSON, N.S.; ZHURINA, F.G.; SENYAVINA, I.B.

Interaction of bromocyanacetic ester with aromatic aldehydes
in the Reformatskii and Witequist reactions. Dokl. AN SSSR 157
no.3:603-606 J1 '64. (MIRA 17:7)

1. Institut khimii prirodnikh soedineniy AN SSSR i Nauchno-
issledovatel'skiy institut organicheskikh poluproduktov i
krasiteley. Predstavleno akademikom M.M. Shemyakinym.

Vul'fson N. S.

75-2-24/64

AUTHORS: Zaretskiy, V. I. , Vul'fson, N. S.

TITLE: Investigations in the Field of Dieckmann's Reaction. (Issledovaniye v oblasti reaktsii Diekmanna) IV. Cyclization of the Tetraethyl-ether of 2-Methylpentantetracarboxylic-1,1,5,5,-Acid (IV. Tsiklizatsiya tetraetilovogo efira 2-metilpentantetrakarbonovoy-1,1,5,5 kisloty)

PERIODICAL: Zhurnal Obshchey Khimii, 1956, Vol. 28, Nr 2, P. 388 - 391 (USSR)

ABSTRACT: The authors continued the investigation of Dieckmann (Diekmann)'s (reference 1) and investigated the cyclization of the tetraethyl-ether of 2-methylpentantetracarboxylic-1,1,5,5-acid (I). It became evident that in the presence of pulverised sodium in benzene the cyclization of tetraether (I) is not possible (the initial tetraether is regenerated). But in the presence of sodium ethylate in an alcohol solution it takes place. 3-methyl-2,6-dicarboethoxy-cyclohexanone (II) forms in a yield of 44,5 %. The cyclization is accompanied by a splitting off of diethylcarbonate (references 2, 3) which is formed during the distillation process of reaction products. The structure of the lacto ether (II) is confirmed by the values of the elementary analysis, of the molecular refraction as well as by the formation of 3-methylcyclohexanone (III), as a re-

Card 1/3

79-2-24/61

Investigations in the Field of Dikman's Reaction. IV. Cyclization of the Tetraethylether of 2-Methylpentantetracarboxylic-1,1,5,5,-Acid

sult of its hydrolysis and the decarboxylation. The constants of the latter correspond to those described in publications for this substance (references 4-6). The initial tetraethylether of 2-methylpentantetracarboxylic-1,1,5,5-acid (I) was obtained by the condensation of 1,3-dibromobutane with sodium diethyl malonate. In the interaction of the stoichiometric quantity of the malonic ester and the dibromide the yield of tetraether (I) amounted to 12,8 %, although sodium bromide was obtained in a quantity of 90 %. In the condensation of 1 Mol of 1,3-dibromobutane with 2 Mol of sodium diethyl malonate and in the presence of a large excess of malonic ester (4 Mol) the tetraether was obtained according to a method analogous to that by V. P. Gol'mov and B. A. Kazanskiy (reference 7), with a yield of 36,9 %. A further increase in the excess of malonic ester does not influence the yield of the condensation product. Conclusions: The cyclization of the tetraethylether of 2-methylpentantetracarboxylic-1,1,5,5-acid was investigated according to Dikman. It was shown that 3-methyl-2,6-dicarboethoxycyclohexane represents the primary reaction product. Its structure was confirmed by the production of 3-methylcyclohexanone with its ketone splitting. There are 11 references, 4 of which are Slavic.

Card 2/3

79-2-24/64

Investigations in the Field of Dikman's Reaction. IV. Cyclization of the Tetra-
ethylether of 2-Methylpentantetracarboxylic-1,1,5,5,-Acid

ASSOCIATION: Scientific Research Institute for Organic Semiproducts and Dyes,
Moscow
(Moskovskiy nauchno-issledovatel'skiy institut organicheskikh polu-
produktov i krasiteley)

SUBMITTED: February 13, 1957

AVAILABLE: Library of Congress

Card 3/3

AUTHORS: Vul'fson, N. S., Zaretskiy, V. I. SOV/79-28-7-41/64

TITLE: Investigation in the Field of Dikman's Reaction (Issledovaniye v oblasti reaktsii Dikmana) V. The Cyclisation of the Diethyl Ester of α -Carbethoxypimelic Acid (V. Tsiklizatsiya dietilovogo efira α -karbetoksipimelinovoy kisloty)

PERIODICAL: Zhurnal obshchey khimii, 1958, Vol. 28, Nr 7, pp. 1909 - 1914 (USSR)

ABSTRACT: The authors investigated the cyclization of the diethyl ester of α -carbethoxypimelic acid according to Dikman (Ref 1). Earlier they had shown that in the cyclization of this ester of α -methylpimelic acid 2 methyl-6-carbethoxycyclohexanone is formed without any doubt. Instead of the normally expected 2-carbethoxycyclohexanone (Formula II) with a simultaneous cleavage of the diethylcarbonate the authors in boiling xylene in the presence of powdery sodium obtained the 2,6-dicarbethoxycyclohexanone (III). It must be noticed that the cyclization of (I) with the same sodium in alcohol solution does practically not take place, while that of the tetraethylesters of the pentane- and

Card 1/3

Investigation in the Field of Dikmar's Reaction. V. SOV/79-28-7-41/64
The Cyclisation of the Diethyl Ester of α -Carbethoxypimelic Acid

2-methyl-pentanetetracarboxylic acids-1,1,5,5 in sodium methylate alcohol solution is accompanied by a cleavage of the diethyl-carbonate and leads to the formation of the β -keto ester (III) (35%) (Ref 3) and 3-methyl-2,6-dicarbethoxycyclohexanone (44%). The formation of (III) can be explained by spatial difficulties. The structure of the compound (III) was proved by its methylation with methyl iodide to (IV) and (V) correspondingly, and after the hydrolysis and decarboxylation of the latter also by the formation of the known cyclohexanones (VI) and (VII). The initial product (I) was produced according to two methods. (Refs 4,5,6). There are 16 references, 5 of which are Soviet.

ASSOCIATION: Nauchno-issledovatel'skiy institut organicheskikh poluproduktov i krasiteley imeni K.Ye.Voroshilova (Scientific Research Institute for Organic Semi-finished Products and Dyes, imeni K. Ye. Voroshilov)

SUBMITTED: May 11, 1956
Card 2/3

Investigation in the Field of Dikmar's Reaction. V. SOV/79-28-7-41/64
The Cyclisation of the Diethyl Ester of α -Carbethoxypimelic Acid

1. Ethyl esters--Chemical reactions 2. Acids--Chemical reactions

Card 3/3

5(3)

AUTHORS: Vinograd, L. Kh., Vul'fson, N. S.

SOV/20-123-1-25/56

TITLE: Reformatskiy's Reaction Involving Nitrobenzaldehydes
(Nitrobenzal'degidy v reaktsii Reformatskogo)

PERIODICAL: Doklady Akademii nauk SSSR, 1958, Vol 123, Nr 1,
pp 97 - 98 (USSR)

ABSTRACT: Any efforts to introduce carbonyl compounds containing
nitro-groups into the Reformatskiy reaction have so
far been a failure (Ref 1). Also the use of these
compounds in the Grignard (Grin'yar) reaction gave
no satisfactory results (Ref 2). The causes of this
failure, however, are different in each of these cases.
In the Grignard reaction the reduction of the nitro-
group by the Grignard reagent is the hindrance. It may
be overcome by a low reaction temperature (Ref 2).
Hence, the nitro-group does not inhibit the reaction
of organometallic compounds with the carbonyl group.
In the Reformatskiy reaction with carbonyl compounds
the total zinc remains practically unchanged. If a

Card 1/3

Reformatskiy's Reaction Involving Nitrobenzaldehydes

SOV/20-123-1-25/56

mixed organozinc compound ought to be formed, in the presence of a carbonyl compound, it can be supposed that the nitro-group contained in it impedes the interaction of zinc with the halogen ester. It was possible to confirm this assumption, since the easily proceeding reaction of acetophenone with bromo acetic ester is completely stopped by addition of 0.05 mol of nitro-benzene. This occurs even if this reaction is carried out in tetrahydrofuran in the presence of mercuric chloride which is known to ease the course of the reaction (Ref 3). Apparently, the Reformatskiy reaction in the presence of carbonyl compounds has to be performed in 2 steps: a) preparation of Reformatskiy reagents from zinc and ester, b) action exercised by the latter on a carbonyl compound. The authors used the method according to reference 4. They succeeded in producing the corresponding esters of β -oxy- β -nitro-phenyl-propionic acids from o-, m- and p-nitro benzaldehydes as well as from bromo acetic ester. The successful carrying out of this reaction confirmed the above assumption, that the nitro group hinders the

Card 2/3

Reformatskiy's Reaction Involving Nitrobenzaldehydes SOV/20-123-1-25/56

formation of mixed organozinc compounds. A characteristic of the resulting products is presented in table 1. There are 1 table and 6 Soviet references.

ASSOCIATION: Nauchno-issledovatel'skiy institut organicheskikh poluproduktov i krasiteley im. K.Ye.Voroshilova (Scientific Research Institute of Organic Semi-Products and Dyes imeni K. Ye. Voroshilov)

PRESENTED: May 23, 1958, by B.A.Kazarskiy, Academician

SUBMITTED: April 16, 1958

Card 3/3

AUTHORS: Vinograd, L. Kh., Vul'fson, H. S. SOV/79-29-1-52/74

TITLE: The Reaction According to Reformatskiy With α -Halogen Nitriles (Reaktsiya Reformatskogo s α -galoidonitrilami)
II. Condensation of Chloro Benzaldehydes With Bromo-Aceto Nitrile (II. Kondensatsiya khlorbental'degidov s bromatseto-nitrilom)

PERIODICAL: Zhurnal obshchey khimii, 1959, Vol 29, Nr 1, pp 245 - 247 (USSR)


ABSTRACT: In the previous reports the authors described the condensation of benzaldehyde with bromo-aceto nitrile according to the reaction by Reformatskiy which proceeds under formation of β -oxy- β -phenyl propionitrile. In continuation of this work they investigated the condensation of the o-, m- and p-chloro benzaldehydes with bromo-aceto nitrile under the usual conditions of this reaction. In this connection they obtained as in the case of application of the non-substituted benzaldehyde the corresponding, previously not described β -oxy- β -chloro-phenyl propionitriles. The nitriles were characterized by the elementary analysis, by the transformation according to Radziszewski (Ref 2)

Card 1/2

The Reaction According to Reformatskiy With α -Halogen Nitriles II. Condensation of Chloro Benzaldehydes With Bromo-Aceto Nitrile

SOV/79-29-1-52/74

into the also hitherto not described oxyamides and by saponification into the well-known trans-o-, m- and p-chlorocinnamic acids: $\text{ClC}_6\text{H}_4\text{CHO} + \text{BrCH}_2\text{CN} \xrightarrow{\text{Zn}} \text{ClC}_6\text{H}_4\text{CH(OH)CH}_2\text{CN}$



CHCOOH . It is worth mentioning that the introduction of the chlorine atom into position 4 of the benzaldehyde increases somewhat the yield in oxynitrile as compared to the non-substituted benzaldehyde (53.4 instead of 48.2%), whereas the introduction of chlorine into position 2, and especially 3 of benzaldehyde leads to a reduction (43.0 and 27%). In the presence of mercury chloride (Ref 3), however, the yield in β -oxy- β -(3-chloro-phenyl)propionitrile increases up to 50.7%. There are 7 references, 3 of which are Soviet.

ASSOCIATION:

SUBMITTED:
Card 2/2

Nauchno-issledovatel'skiy institut poluproduktov i krasiteley (Scientific Research Institute of Intermediate Products and Dyes)
November 29, 1957

AUTHORS: Zaretskiy, V. I., Vul'fson, N. S. SOV/79-29-2-13/71

TITLE: Investigation in the Field of the Dikman Reaction (Issledovaniye v oblasti reaktsii Dikmana). VI. Cyclization of Diethyl Ester of α -Acetyl and α -Benzoyl Pimelic Acid (VI. Tsiklizatsiya dietilovogo efira α -atsetil- i α -benzoilpimelinovoy kisloty)

PERIODICAL: Zhurnal obshchey khimii, 1959, Vol 29, Nr 2, pp 416-421 (USSR)

ABSTRACT: The cyclization of α -acetyl (I) and α -benzoyldiethyl pimelate (II) according to Dikman is investigated here. The authors had already earlier shown that the cyclization of α -carbethoxydiethyl pimelate takes place unexpectedly on the non-substituted α -carbon atom and leads to 2,6-dicarbethoxycyclohexanone. It was of interest to clarify whether the cyclization (I) takes place on the substituted α -carbon atom in the direction to the formation of 2-carbethoxycyclo-hexanone (III) (with the cleavage of ethyl acetate) or of 2-acetylcyclo-hexanone (IV) (with the cleavage of diethyl carbonate) (Scheme 1). Actually, in the cyclization of (I) in boiling xylene in the presence of 1.4 g-at. powdery sodium or 1.4 mol sodium ethylate, compound (III) in 52-57% yield is formed, in which connection ethyl acetate separates. In the reaction of 1.5 mol sodium ethylate in alcohol

Card 1/3

SOV/79-29-2-13/71

Investigation in the Field of the Dikman Reaction. VI. Cyclization of Diethyl Ester of α -Acetyl and α -Benzoyl Pimelic Acid

solution, diethyl pimelate formed as chief product (V) in a yield of 35%, whereas the yield in (III), which separated in form of 2-phenyl-4,5,6,7-tetrahydro-indazoline-3, amounted to only 6.5%. Compound (III) cleaves into (V) only to a small extent (3.1%), whereas (I), on boiling with the alcohol solution of 0.2 mol sodium ethylate it forms compound (V) (2.2% yield), and with 1 mol sodium ethylate it yields 56%. The formation of small quantities (III), on processing (I) with the alcohol solution of 1.5 mol sodium ethylate, can be explained by the cyclization of (V) which takes place in a low yield. The problem concerning the reaction mechanism in the presence of powdery sodium was solved with smaller quantities of sodium (0.9 g-at). Ethylacetate (27.3%), (V, 13.9%) and (III, 16%) proved to be the chief products in this connection. Thus, alcoholysis (I) appears as the first process stage in non-alcoholic medium, under formation of compound (V), which then cyclizes into (III). An interpretation of this alcoholysis is suggested and the investigation results obtained are used to set up a scheme of the cyclization mechanism of α -acetyl and α -benzoyldiethyl pimelate (Scheme 2).

Card 2/3

SOV/79-29-2-13/71

Investigation in the Field of the Dikman Reaction. VI. Cyclization of Diethyl Ester of α -Acetyl and α -Benzoyl Pimelic Acid

The structure of 2-carboethoxycyclohexanone (III) was confirmed by the synthesis of the known 2-phenyl-4,5,6,7-tetrahydro-indazolone-3 from cyclohexanone. There are 11 references, 6 of which are Soviet.

ASSOCIATION: Nauchno-issledovatel'skiy institut organicheskikh poluproduktov i krasiteley (Scientific Research Institute for Organic Semi-products and Dyes)

SUBMITTED: December 31, 1957

Card 3/3

5 (3)

AUTHORS:

Vul'fson, N. S. Vinograd, L. Kh.

SCY/79-29-1-24/77

TITLE:

Reformatskiy's Reaction With α -halogen Nitriles (Reaktsiya Reformatskogo s α -galojdnitrilami). III. Condensation of the Methoxy-benzaldehydes With Bromo-aceto Nitrile (III. Kondensatsiya metoksibenzal'dogidov s bromoacetonnitrilom)

PERIODICAL:

Zhurnal obshchey khimii, 1959, Vol 29, Nr 4, pp 1147-1149 (USSR)

ABSTRACT:

The authors previously described the condensation of benzaldehyde and chloro-benzaldehydes with bromo-aceto nitrile (Refs 1, 2). Further, the condensation of bromo-aceto nitrile with o-, m-, p-methoxy- and 3,4-dimethoxy-benzaldehydes were investigated under the usual conditions of Reformatskiy's reaction. The β -(2-methoxy-phenyl)- and β -(3-methoxy-phenyl)- β -oxy-propionitriles, which so far have not yet been described, were accordingly obtained from o- and m-methoxy-benzaldehydes as well as with chlorobenzaldehydes and unsubstituted benzaldehyde. In the case of p-methoxy- and 3,4-dimethoxy-benzaldehydes the known nitriles of p-methoxy- and 3,4-dimethoxy-cinnamic acids were formed (Refs 3, 4). The β -(2-methoxy-phenyl)- and β -(3-methoxy-phenyl)- β -oxy-

Card 1/3
2

Reformatskiy's Reaction With α -Halogen Nitriles.

SOV/79-29-4-24/77

III. Condensation of the Methoxy-benzaldehydes With Bromo-aceto Nitrile

propionitrile were transformed by hydrogenperoxide into the amides of β -(2-methoxy-phenyl)- and β -(3-methoxy-phenyl)- β -oxy-propionic acids, and by saponification into the 2- and 3-methoxy-cinnamic acids. The condensation of m-methoxy-benzaldehyde and m-chloro-benzaldehyde (Ref 2) was carried out in a lower yield than with the o- and p-isomers, while mercury chloride increased the yield only from 22.1 to 25.7 %. Very good yields were attained with tetrahydrofuran instead of the usual solvents (Ref 5). In this connection the yields for o-, m-, p-methoxy- and 3,4-dimethoxy-benzaldehydes were increased from 49.7 to 70.4 %, from 25.7 to 74 %, from 34.6 to 62 %, and from 23 to 76.3 %, accordingly. The reaction in tetrahydrofuran proceeded abruptly, thus shortening reaction time and decreasing resin formation. There are 10 references, 4 of which are Soviet.

ASSOCIATION:

Nauchno-issledovatel'skiy institut organicheskikh poluproduktov i krasiteley (Scientific Research Institute of Organic Semi-products and Dyes)

Card 2/3

2

KOROLEV, A.I., otv.red.; VUL'FSON, N.S., zam.otv.red.; BOGDANOV, S.V., red.; DOKUNIKHIN, N.S., red.; MASLENNIKOVA, Ye.V., red.; PODIMAN, I.V., red.; KHOMSKIY, I.G., red.; ZHITIN, V.I., red.; SHPAK, Ye.G., tekhn.red.

[Organic intermediate products and dyes; collected articles]
Organicheskie poluprodukty i krasiteli; sbornik statei. Moskva,
Gos.nauchno-tekhn.izd-vo khim.lit-ry. No.1. 1959. 238 p.
(MIRA 13:7)

1. Nauchno-issledovatel'skiy institut organicheskikh poluproduktov i krasiteley.

(Dyes and dyeing)

(Aromatic compounds)

VUL'FSON, N.S.; IODKO, M.O.

Investigations in the field of Dieckmann reaction. Report
No.1: Synthesis of α -phenylene acetic- β -propionic acid and
its ester tagged with C^{14} in the carboxylic group of the
propionic acid residue. Org. poluprod. i kras. no.1:92-
95 '59. (MIRA 14:11)

(Propionic acid)
(Radioactive tracers)
(Dieckmann condensation)

STEPANOV, F. N.; VUL'FSON, N. S.

Investigation in the series of derivatives of acetonitrile. Report
No.1. Hydrolytic splitting of acrylcynoacetic esters. Org.
poluprod. i kras. no.1:222-230 '59. (MIRA 14:11)
(Acetonitrile)
(Cyanoacetic acid)

VUL'FSON, N. S.; DAVIDOVA, S. L.; LUKASHINA, L. I.

Investigation in the series of derivatives of acetonitrile. Report
No.2: 2-cyanomethylquinoxalone-3 and its derivatives. Org. polu-
prod. 1 kras. no.1:231-236 '59. (MIRA 14:10)
(Acetonitrile)

AUTHORS: Vinograd, L. Kh., Vul'fson, N. S. SOV/79-29-8-53/31

TITLE: Reformatskiy's Reaction With α -Halogen Nitriles. IV. Condensation of Ketones With Bromoacetonitrile

PERIODICAL: Zhurnal obshchey khimii, 1959, Vol 29, Nr 8, pp 2690-2692 (USSR)

ABSTRACT: In previous papers (Ref 1), the reaction of bromoacetonitrile with different aromatic aldehydes was described which takes place according to Reformatskiy under usual conditions, and yields the β -oxynitriles. The authors carried on with the investigation of this reaction, and condensed ketones (acetophenone, cyclohexanone and dibutyl ketone) with bromoacetonitrile; the corresponding β -oxynitriles resulted. On distillation of β -oxy- β -phenylbutyronitrile, a partial dehydration takes place, and an impurity of the nitrile of the β -methyl-cinnamic acid is formed. The oxynitrile can be isolated from the higher-boiling fraction by crystallization. The pure, unsaturated nitrile was obtained by dehydration of the nitrile mixture with potassium bisulfate. The solvent used influences the course of reaction. In the case of the reaction of bromoacetonitrile with acetophenone, it was found that in benzene, toluene and in a mixture of benzene and ether mainly resinous products are formed; in dioxane, ether and their

Card 1/2

Reformatskiy's Reaction With α -Halogen Nitriles. IV.
Condensation of Ketones With Bromoacetonitrile

SOV/79-29-8-53/31

mixture the maximum yields are 38.5%; the best yields (67.8%) were obtained when using tetrahydrofuran (Table 1). The nitriles not described in publications, with the exception of 5-oxy-5-cyanomethylnonane, were converted into the corresponding amides and, besides, the nitrile of the β -methyl-cinnamic acid was converted into the β -methyl-cinnamic acid. Thus, the nitriles of the β -methyl-cinnamic-, β -oxy- β -phenyl-butyric-, and β -oxy- β -butyl-heptanic acid were synthesized. The results of the experiments are given in table 2, the characteristic features of the amides in table 3. There are 3 tables and 3 references, 2 of which are Soviet.

ASSOCIATION: Nauchno-issledovatel'skiy institut organicheskikh poluproduktov i krasiteley (Scientific Research Institute for Organic Semi-products and Dyes)

SUBMITTED: July 11, 1958

Card 2/2

3(3)

SOV/79-29-8-54/81

AUTHORS: Vul'fson, N. S., Vinograd, L. Zh.

TITLE: Reformatskiy's Reaction With α -Halogen Nitriles. V. Reaction of Benzaldehyde and Acetophenone With Chloroaceto-, α -Bromopropio-, and α -Bromoisobutyro Nitriles

PERIODICAL: Zhurnal obshchey khimii, 1959, Vol 29, Nr 8, pp 2692-2695 (USSR)

ABSTRACT: The authors tried to extend Reformatskiy's reaction with bromoacetonitrile described previously (Ref 1) to other α -halogen nitriles: to chloroacetonitrile, α -bromopropionitrile, and α -bromoisobutyronitrile. All these nitriles were allowed to react with benzaldehyde and acetophenone. In analogy with the reactions of chloroacetate, the chloroacetonitrile reacts more difficultly than the corresponding bromide, as was expected, and gives smaller yields in β -oxynitriles. The α -methyl derivatives of bromoacetonitrile (α -bromopropionitrile and α -bromoisobutyronitrile) react more readily than bromoacetonitrile and give better yields than the corresponding β -oxynitriles. The nitriles were identified, as previously (Ref 1), by transformation into the corresponding β -oxyamides. Yet, not all nitriles reacted in this way. The reaction time of α -methyl- and α,α -dimethyl- β -oxy- β -phenylpropionitrile with H_2O_2 had to be prolonged to 3 hours, and that of

Card 1/2

Reformatskiy's Reaction With α -Halogen Nitriles. V. SOV/79-29-6-54/81
 Reaction of Benzaldehyde and Acetophenone With Chloroaceto-, α -Bromopropio-,
 and α -Bromoisobutyro Nitriles

α -methyl- and α,α -dimethyl- β -oxy- β -phenylbutyronitrile to 6 hours. Only the first three nitriles gave the corresponding amides in small yields. In the case of α,α -dimethyl- β -oxy- β -phenylbutyronitrile, even a splitting-up of the carbon chain takes place, and acetophenone is formed again. On hydrolysis of nitriles (boiling with KOH for 12-15 hours) surprisingly no cinnamic acids resulted. This can be explained by the influence exerted by the α -substituents upon the stability of the carbon chain. The experimental results are given in two tables. There are 2 tables, and 11 references, 3 of which are Soviet.

ASSOCIATION: Nauchno-issledovatel'skiy institut organicheskikh poluproduktov i krasiteley (Scientific Research Institute for Organic Semi-products and Dyes)

SUBMITTED: July 11, 1958

Card 2/2

5(3).

AUTHORS: Vul'fon, N. S., Zaretskiy, V. I.

SOV/79-29-8-64/81

TITLE: Investigations in the Field of the Dickmann Ring Formation.
VII. Regrouping of 2-methyl-2-carbethoxycyclohexanone Forming
6-methyl-2-carbethoxycyclohexanone

PERIODICAL: Zhurnal obshchey khimii, 1959, Vol 29, Nr 8, pp 2737-2738 (USSR)

ABSTRACT: Since the publications confine themselves (Refs 1-5) to the description of the regrouping of the 2-carbethoxyalkyl derivatives of the 2-carbethoxycyclohexanone, it was interesting for the authors to investigate the analogous regrouping of the 2-alkyl derivative starting from 2-methyl-2-carbethoxycyclohexanone (I). As it is known, the diethyl- α -methylpimelinate (Ref 6) the cyclization of which in boiling benzene in the presence of powdered sodium according to Dickmann yields, as previously reported (Ref 7), the compound (II), is easily formed when compound (I) is boiled in an alcohol solution of sodium ethylate (0,2 mol). However, under usual conditions diethyl- α -methylpimelinate (40%), the unchanged β -keto ester (I) (20%), and traces only of the β -keto ester (II) were obtained, whereas the boiling of 2-methyl-2-carbethoxycyclohexanone in xylene in the

Card 1/2

Investigations in the Field of the Dickmann Ring SOV/79-29-8-64/81
Formation. VII. Regrouping of 2-methyl-2-carbethoxycyclohexanone Forming
6-methyl-2-carbethoxycyclohexanone

presence of sodium ethylate yields the regrouping forming compound (II) with a 42% yield (Scheme). It was found that diethyl- α -methylpimelinate does not cyclize in the alcohol solution of sodium ethylate, while its ring formation in boiling xylene in the presence of sodium ethylate proceeds smoothly and yields the compound (II) (54%). These facts as well as the formation of a large amount of α -methylpimelinate in the course of the attempt to carry out the regrouping of (I) in an alcohol medium which failed are convincing proof that the regrouping of (I) into the analogous 2,6-compound takes place by an opening of the ring and subsequent cyclization of the resulting diethyl- α -methylpimelinate. There are 8 references, 1 of which is Soviet.

ASSOCIATION: Nauchno-issledovatel'skiy institut organicheskikh poluproduktov i krasiteley imeni K. Ye. Voroshilova (Scientific Research Institute of Organic Semi-finished Products and Dyes imeni K. Ye. Voroshilov)

SUBMITTED: July 4, 1958

Card 2/2

V. L'ISOI, I.S.; ZHUKOV, K.S.

Synthesis of 3-acyl-4-hydroxy-carbostyrils. Zhur. TSO 5
no. 3:352-353 '60. (I.M. 4:2)

1. Nauchno-issledovatel'skiy institut organicheskikh poluproduktov
i koksitolov Irini K.Ye. Voroshilova.
(Carbostyryl)

ZHURIN, R.B.; VULFSON, N.S.

Reaction of C-acyla ion of heterocyclic ketoenols. Part 2: Synthesis of α -acetyl- and α -propionyl- γ -phenyltetronic acids. Zhur.ob.khim. 30 no.8:2467-2468 Ag '60. (MIRA 13:8)

1. Nauchno-issledovatel'skiy institut organicheskikh poluproduktov i krasiteley imeni K.Ye.Voroshilova.
(Tetronic acid)

KOLCHIN, V.Ye.; VUL'FSON, N.S.

Derivatives of acylacetic esters of the heterocyclic series. Part 2:
Arylides of α - and β -furoylacetic acids, and synthesis of azomethine
dyes from them. Zhur. ob. khim. 30 no.9:3091-3095 S '60.

(MIRA 13:9)

(Furanpropionic acid)

(Dyes and dyeing)

VUL'FSON, N.S.

Investigation in the series of derivatives of acetonitrile. Report
No. 3: Synthesis of substituted ω -cyanoacetophenones. Org. polu-
prod. i kras. no.2:128-136 '61. (MIRA 14:11)
(Acetophenone) (Acetonitrile)

VUL'FSON, N.S.; IODKO, M.O.

Synthesis of β -tetralone. Org. poluprod. i kras. no.2:143-145
'61. (MIRA 14:11)
(Naphthalenone)

VUL'FSON, N.S.; ZHURIN, R.B.

APPROVED FOR RELEASE: 09/01/2001: 3-CIA-RDP86-00513R001961310009-1"

Cyclization of phenylhydrazones of 3-acetyl-4-hydroxy-2-pyridone
Zhur. VKHO 6 no.2:239-240 '61. (MIRA 14:3)

1. Nauchno-issledovatel'skiy institut organicheskikh poluproduktov
i krasiteley imeni K. Ye. Voroshilova.
(Hydrazones) (Coumarin)

VUL'FSON, N.S.; KOLCHIN, V.Ye.

Derivatives of acylacetic esters of the heterocyclic series. Part 3:
Synthesis of α - and β -thenoyl acetic esters and arylides, and of
azomethine dyes based on them. Zhur.ob.khim. 30 no.10:3425-3430
O '61. (MIRA 14:4)

1. Nauchno-issledovatel'skiy institut organicheskikh poluproduktov
i krasiteley.
(Dyes and dyeing) (Thiophenecarboxylic acid)

VUL'FSON, N.S.; ZHURIN, R.B.

Reactions of C-acylation of heterocyclic keto-enols. Part 4:5-Acylbarbituric acids. Zhur. ob. khim, 31 no.1:261-283 Ja '61.

(MIRA 14:1)

1. Nauchno-issledovatel'skiy institut organicheskikh poluproduktov i krasiteley.

(Barbituric acid)

ZARETSKIY, V.I.; VUL'FSON, N. S.

Dieckman reactions. Part 8: Cyclization of the diethyl
ester of α -ethyl- α -carbethoxypimelic acid. Zhur. ob. khim.
31 no. 2:484-490 F '61. (MIRA 14:2)

1. Institut khimii prirodnikh soedineniy AN SSSR.
(Heptanedioic acid) (Cyclization)

ZHURIN, R.B.; VUL'FSON, N.S.

C-Acylation of heterocyclic ketoenols. Part 5: Mechanism of
the C-acylation of 4-hydroxycoumarin. Zhur. ob. khim, 31
no.3:875-879 Mr '61. (MIRA 14:3)

1. Institut organicheskikh poluproduktov i krasiteley imeni
K. Ye. Voroshilova.
(Coumarin) (Acylation)

VUL'FSON, N.S.; ZHURIN, R.B.

C-acylation of heterocyclic keto ~~enols~~. Part 6: Cyclization
of phenylhydrazones of 3-acyl-4-hydroxycoumarins. Zhur.ob.khim.
31 no.10:3381-3385 0 '61. (MIRA 14:10)

1. Institut organicheskikh poluproduktov i krasiteley imeni
K.Ye.Voroshilova.
(Coumarin) (Hydrazones)

VUL'FSON, N.S.

10

Probably
N.M.

SEVERIN, Sergey Yevgen'yevich, Institute of Pharmacology and Chemotherapy, Academy of Medical Sciences, Moscow; VUL'FSON, N. S. [possibly P.L. VUL'FSON, Chair, Animal Biochemistry, Moscow State University (1959 position)] - "The importance of karnosis in neurotrophic relations" Session I

SHAMARINA, N. N., Physiological Laboratory, Academy of Sciences USSR, Moscow - "Effect of tetanic stimulation on different muscle fibers" II-2-b

STUDITSKIY, Aleksandr Nikolayevich, ZHEREVSKAYA, R. P., and RUMYANTSEVA, O.N., all of the Institute of Animal Morphology Lenin A. N. Severtsov, Academy of Sciences USSR, Moscow - "Neurotrophic influence in recovery of structure and function of regenerating muscle" I

TELEPIEVA, V. I., Chair, Animal Biochemistry, Moscow State University, Moscow - "Changes in muscle following denervation" Session II-2-a

YAKOVLEV, N. N., KRASHOVA, A. F., and CHAGOVETS, N.R., all of the Leningrad Scientific Research Institute, Institute of Physical Culture, Leningrad - "Adaptation of energy metabolism in muscle" Session II-2-b

Report to be submitted for the Symposium on the Effects of Use and Disease on Neuromuscular Functions (IUPS), Prague-Liblice, Czech, 18-24 Sep 1962.

VUL'FSON, N.S.; ZARETSKIY, V.I.; DOMANINA, O.N.

Synthesis of ethyl- β -carbethoxyethyl ester of malonic
acid. Zhur. VKHO 7 no.6:709 '62. (MIRA 15:12)

1. Institut khimii prirodnikh soedineniy AN SSSR.
(Malonic acid)

VUL'FSON, N.S.; ZHURIN, R.B.

Reaction of cycloacylation of heterocyclic keto-enols. Part 7:
Cyclization of phenyl-hydrazones of 3-acyl-4-hydroxycarbostryls.
Zhur.ob.khim. 32 no.3:991-994 Mr '62. (MIRA 15:3)

1. Institut organicheskikh poluproduktov i krasiteley i Institut
khimii prirodnykh soyedineniy AN SSSR.
(Hydrazones) (Carbostryls) (Cyclization)

VUL'FSON, N.S.; PODREZOVA, T.N.

Dieckmann condensation. Part 9: Cyclization of ethyl ester of
o-(carbethoxymethoxy) phenylacetic acid. Zhur.ob.khim. 32
no.9:3019-3022 S '62. (MIRA 15:9)

1. Institut khimii prirodnikh soedineniy AN SSSR i Nauchno-
issledovatel'skiy institut organicheskikh poluproduktov i
krasiteley.

(Acetic acid) (Dieckmann condensation)

VUL'FSON, N.S.; KOLCHIN, V.Ye.; ARTEMCHIK, L.K.

Derivatives of acylacetic esters of the heterocyclic series. Part 4: Synthesis of nicotinoylacetic ester, arylides, and azomethine dyes prepared from them. Zhur.ob.khim. 32 no.10:3382-3386 0 '62. (MIRA 15:11)

1. Nauchno-issledovatel'skiy institut organicheskikh poluproduktov i krasiteley i Institut khimii prirodnykh soedineniy AN SSSR.

(Pyridinepropionic acid)
(Dyes and dyeing) (Schiff bases)